Towards Metric Temporal Answer Set Programming

PEDRO CABALAR*

University of Corunna, Spain

MARTÍN DIÉGUEZ

LERIA, Université d'Angers, France

TORSTEN SCHAUB and ANNA SCHUHMANN

University of Potsdam, Germany

submitted 1 January 2003; revised 1 January 2003; accepted 1 January 2003

Abstract

We elaborate upon the theoretical foundations of a metric temporal extension of Answer Set Programming. In analogy to previous extensions of ASP with constructs from Linear Temporal and Dynamic Logic, we accomplish this in the setting of the logic of Here-and-There and its non-monotonic extension, called Equilibrium Logic. More precisely, we develop our logic on the same semantic underpinnings as its predecessors and thus use a simple time domain of bounded time steps. This allows us to compare all variants in a uniform framework and ultimately combine them in a common implementation.

This article is under consideration for acceptance in TPLP.

1 Introduction

Reasoning about action and change, or more generally reasoning about dynamic systems. is not only central to knowledge representation and reasoning but at the heart of computer science. We addressed this over the last years by combining traditional approaches, like Dynamic and Linear Temporal Logic (DL (Harel et al. 2000) and LTL (Pnueli 1977)), with the base logic of Answer Set Programming (ASP (Lifschitz 1999)), namely, the logic of Here-and-There (HT (Heyting 1930)) and its non-monotonic extension, called Equilibrium Logic (EL (Pearce 1997). This resulted in non-monotonic linear dynamic and temporal equilibrium logics (DEL (Bosser et al. 2018; Cabalar et al. 2019) and TEL (Aguado et al. 2013; Cabalar et al. 2018)) that gave rise to the temporal ASP system telingo (Cabalar et al. 2019; Cabalar et al. 2020) extending the full-featured ASP system clingo (Gebser et al. 2016). A key design decision has been to base both logics on the same semantic structures so that language constructs from both can be jointly used in an implementation. Another commonality of dynamic and temporal logics is that they abstract from specific time points and rather focus on capturing temporal relationships. For instance, we can express in a temporal logic that a machine has to be eventually cleaned after being used with the formula $\Box(use \rightarrow \Diamond clean)$. However, sometimes this is not enough to capture the desired relation. That is, we might also want to quantify the time difference between both

^{*} Partially supported by MINECO, Spain, grant TIC2017-84453-P.

events. For instance, whenever the machine is used, it has to be cleaned within less than 5 time units. This can be expressed by means of metric temporal operators as follows:

$$\Box(use \to \Diamond_5 clean) \ . \tag{1}$$

In this paper, we address this issue and elaborate upon a combination of Metric Temporal Logic (MTL (Alur and Henzinger 1992; Ouaknine and Worrell 2005)¹) with HT and EL. Our development of *Metric Equilibrium Logic* (MEL) not only parallels the one of TEL and DEL mentioned above but, moreover, builds on the same semantic foundations. This allows us to relate all three systems in a uniform semantic setting and, ultimately, to integrate the corresponding language constructs in a common implementation.

A full version of this paper including proofs of results can be found at http://arxiv.org/abs/2008.02038.

2 Metric Equilibrium Logic

Given a set \mathcal{A} of atoms, or *alphabet*, we define a *(metric)* formula φ by the grammar:

$$\varphi ::= a \mid \perp \mid \varphi_1 \bowtie \varphi_2 \mid \bullet \varphi \mid \varphi_1 \mathbf{S}_n \varphi_2 \mid \varphi_1 \mathbf{T}_n \varphi_2 \mid \circ \varphi \mid \varphi_1 \mathbb{U}_n \varphi_2 \mid \varphi_1 \mathbb{R}_n \varphi_2$$

where $a \in \mathcal{A}$ is an atom and $\bowtie \in \{\rightarrow, \land, \lor\}$ is a binary Boolean connective; n is a numeral constant (referring to some integer number) or the symbolic constant ℓ (standing for the length of a trace; see below). The last six cases of φ correspond to the metric past connectives *previous*, *since*, *trigger*, and their future counterparts *next*, *until*, and *release*, where n > 0 restricts the scope of each operator to the last (resp. next) n time points, including the current state.²

We also define several derived operators like the Boolean connectives $\top \stackrel{def}{=} \neg \bot$, $\neg \varphi \stackrel{def}{=} \varphi \rightarrow \bot$, $\varphi \leftrightarrow \psi \stackrel{def}{=} (\varphi \rightarrow \psi) \land (\psi \rightarrow \varphi)$, and the following derived metric operators:

$$\begin{array}{cccc} \mathbf{I} & \stackrel{def}{=} & \neg \bullet \top & \mathbb{F} & \stackrel{def}{=} & \neg \bullet \top \\ \widehat{\bullet}\varphi & \stackrel{def}{=} & \bullet \varphi \lor \mathbf{I} & \widehat{\circ}\varphi & \stackrel{def}{=} & \circ \varphi \lor \mathbb{F} \\ \mathbf{I}_n \varphi & \stackrel{def}{=} & \bot \mathbf{T}_n \varphi & \Box_n \varphi & \stackrel{def}{=} & \bot \mathbb{R}_n \varphi \\ \mathbf{\diamond}_n \varphi & \stackrel{def}{=} & \top \mathbf{S}_n \varphi & \Diamond_n \varphi & \stackrel{def}{=} & \top \mathbb{U}_n \varphi \end{array}$$

On the left, we give past operators, namely, *initial, weak previous, always before, eventually before*, while the right column lists their future counterparts *final, weak next, always afterward, eventually afterward.* We define the iterated application of the one step operators as $\otimes^0 \varphi \stackrel{def}{=} \varphi$ and $\otimes^n \varphi \stackrel{def}{=} \otimes \otimes^{n-1} \varphi$ for n > 0 and $\otimes \in \{\bullet, \circ, \widehat{\bullet}, \widehat{\circ}\}$. For instance, $\circ^2 p$ corresponds to $\circ \circ p$. Here, the use of n with temporal operators captures a number of iterations of some one step expression so that, as we see below, for instance, $\diamond_3 \varphi$ amounts to $\varphi \vee \circ \varphi \vee \circ^2 \varphi$.

An example of metric formulas is the modeling of traffic lights. While the light is red by default, it changes to green within less than 3 time units whenever the button is pushed; and it stays green for other 3 time units. This can be represented by

$$\Box(red \land green \to \bot) \tag{2}$$

¹ Unlike traditional approaches usually having continuous time domains (Alur and Henzinger 1992), we deal with point-based semantics based on discrete linear time, similar to (Ouaknine and Worrell 2005).

² Values $n \leq 0$ are tolerated but trivialize the subformula at hand, as made precise in Proposition 6.

$$\Box(\neg green \to red) \tag{3}$$

$$\Box(push \to \Diamond_3 \Box_4 green) \tag{4}$$

Note that this example combines a default rule (3) with a metric rule (4), describing the initiation and duration period of events. This nicely illustrates the interest in nonmonotonic metric representation and reasoning methods.

Given $a \in \mathbb{N}$ and $b \in \mathbb{N} \cup \{\omega\}$, we let [a..b] stand for the set $\{i \in \mathbb{N} \mid a \leq i \leq b\}$, [a..b)for $\{i \in \mathbb{N} \mid a \leq i < b\}$ and (a..b] for $\{i \in \mathbb{N} \mid a < i \leq b\}$. For the semantics, we start by defining a *trace* of length λ over alphabet \mathcal{A} as a sequence $(H_i)_{i \in [0..\lambda)}$ of sets $H_i \subseteq \mathcal{A}$. A trace is *infinite* if $\lambda = \omega$ and *finite* if $\lambda = n$ for some natural number $n \in \mathbb{N}$. Given traces $\mathbf{H} = (H_i)_{i \in [0..\lambda)}$ and $\mathbf{H}' = (H'_i)_{i \in [0..\lambda)}$ both of length λ , we write $\mathbf{H} \leq \mathbf{H}'$ if $H_i \subseteq H'_i$ for each $i \in [0..\lambda)$; accordingly, $\mathbf{H} < \mathbf{H}'$ iff both $\mathbf{H} \leq \mathbf{H}'$ and $\mathbf{H} \neq \mathbf{H}'$.

Our semantics is based on *Here-and-There traces* (for short HT-*traces* (Cabalar et al. 2018)) of length λ over alphabet \mathcal{A} being sequences of pairs $(\langle H_i, T_i \rangle)_{i \in [0.\lambda)}$ such that $H_i \subseteq T_i \subseteq \mathcal{A}$ for any $i \in [0.\lambda)$. We often represent an HT-trace as a pair of traces $\langle \mathbf{H}, \mathbf{T} \rangle$ of length λ where $\mathbf{H} = (H_i)_{i \in [0.\lambda)}$ and $\mathbf{T} = (T_i)_{i \in [0.\lambda)}$ such that $\mathbf{H} \leq \mathbf{T}$. When an HT-trace $\langle \mathbf{H}, \mathbf{T} \rangle$ satisfies $\mathbf{H} = \mathbf{T}$, it is called *total*.

We assume a one-to-one correspondence between numeral constants and integers and let \overline{n} stand for the number corresponding to numeral n. For the symbolic constant ℓ , we fix $\overline{\ell} = \lambda$ to the length λ of the trace. For simplicity, we let expressions like n - m, formed with numeral constants n and m, stand for the numeral representing the difference between \overline{n} and \overline{m} .

We define the semantics of metric formulas in terms of HT-traces.

Definition 1 (Satisfaction)

Let $\mathbf{M} = \langle \mathbf{H}, \mathbf{T} \rangle$ be an HT-trace of length λ over alphabet \mathcal{A} , and let φ be a metric formula over \mathcal{A} . The trace \mathbf{M} satisfies φ at time point $k \in [0..\lambda)$, written $\mathbf{M}, k \models \varphi$, if

- 1. $\mathbf{M}, k \not\models \bot$
- 2. $\mathbf{M}, k \models a \text{ iff } a \in H_k$, for any atom $a \in \mathcal{A}$
- 3. $\mathbf{M}, k \models \varphi \land \psi$ iff $\mathbf{M}, k \models \varphi$ and $\mathbf{M}, k \models \psi$
- 4. $\mathbf{M}, k \models \varphi \lor \psi$ iff $\mathbf{M}, k \models \varphi$ or $\mathbf{M}, k \models \psi$
- 5. $\mathbf{M}, k \models \varphi \rightarrow \psi$ iff $\langle \mathbf{H}', \mathbf{T} \rangle, k \not\models \varphi$ or $\langle \mathbf{H}', \mathbf{T} \rangle, k \models \psi$, for all $\mathbf{H}' \in \{\mathbf{H}, \mathbf{T}\}$
- 6. $\mathbf{M}, k \models \mathbf{\Phi}\varphi$ iff k > 0 and $\mathbf{M}, k-1 \models \varphi$
- 7. $\mathbf{M}, k \models \circ \varphi$ iff $k + 1 < \lambda$ and $\mathbf{M}, k + 1 \models \varphi$
- 8. $\mathbf{M}, k \models \varphi \bigcup_n \psi$ iff for some $j \in [0..\overline{n})$ such that $k+j \in [0..\lambda)$ we have $\mathbf{M}, k+j \models \psi$ and $\mathbf{M}, k+i \models \varphi$ for all $i \in [0..j)$
- 9. $\mathbf{M}, k \models \varphi \mathbb{R}_n \psi$ iff for all $j \in [0..\overline{n})$ such that $k+j \in [0..\lambda)$ we have $\mathbf{M}, k+j \models \psi$ or $\mathbf{M}, k+i \models \varphi$ for some $i \in [0..j)$
- 10. $\mathbf{M}, k \models \varphi \mathbf{S}_n \psi$ iff for some $j \in [0..\overline{n})$ such that $k-j \in [0..\lambda)$ we have $\mathbf{M}, k-j \models \psi$ and $\mathbf{M}, k-i \models \varphi$ for all $i \in [0..j)$
- 11. $\mathbf{M}, k \models \varphi \mathbf{T}_n \psi$ iff for all $j \in [0..\overline{n})$ such that $k-j \in [0..\lambda)$ we have $\mathbf{M}, k-j \models \psi$ or $\mathbf{M}, k-i \models \varphi$ for some $i \in [0..j)$

The fundamental difference to standard temporal logics is clearly the satisfaction of implication ' \rightarrow ' that is inherited from the (non-temporal) logic HT, an intermediate logic dealing with exactly two worlds h, t with the reflexive accessibility relation $h \leq t$. When traces are total $\langle \mathbf{T}, \mathbf{T} \rangle$, we get h = t and this distinction disappears, so implication

becomes classical. From the perspective of metric temporal logic, the definition of release and trigger in 9 and 11, respectively, are additionally conditioned by the trace's limits and can thus be seen as weak variants of the standard counterparts. Similarly, the satisfaction of until and since formulas in 8 and 10, respectively, is restricted to the time points within a trace. Clearly, for infinite traces, the restriction of future operators vanishes.

An HT-trace **M** is a *model* of a metric theory Γ if $\mathbf{M}, 0 \models \varphi$ for all $\varphi \in \Gamma$. A formula φ is a *tautology* (or is *valid*), written $\models \varphi$, iff $\mathbf{M}, k \models \varphi$ for any HT-trace **M** and any $k \in [0..\lambda)$. We call the logic induced by the set of all tautologies *Metric logic of Here* and *There* (MHT for short). We say that an HT-trace **M** is a *model* of a set of formulas (or *theory*) Γ iff $\mathbf{M}, 0 \models \varphi$ for any $\varphi \in \Gamma$. Two formulas φ, ψ are *equivalent* if $\models \varphi \leftrightarrow \psi$. Whenever two formulas φ and ψ are equivalent, they are completely interchangeable in any theory without altering the theory's semantics.

We write $\operatorname{MHT}(\Gamma, \lambda)$ to stand for the set of models of length λ of a theory Γ , and define $\operatorname{MHT}(\Gamma) \stackrel{def}{=} \operatorname{MHT}(\Gamma, \omega) \cup \bigcup_{\lambda \geq 0} \operatorname{MHT}(\Gamma, \lambda)$, that is, the whole set of models of Γ of any length. An interesting subset of $\operatorname{MHT}(\Gamma, \lambda)$ is the one formed by total traces $\langle \mathbf{T}, \mathbf{T} \rangle$, we denote as $\operatorname{MTL}(\Gamma, \lambda)$. We also use $\operatorname{MTL}(\Gamma)$ to stand for $\operatorname{MHT}(\Gamma, \omega) \cup \bigcup_{\lambda \geq 0} \operatorname{MTL}(\Gamma, \lambda)$. In the non-metric version of temporal HT, the restriction to total models turns out to correspond to Linear Temporal Logic (LTL). In our case, it defines a metric version of LTL that we call *Metric Temporal Logic* (MTL for short). It can be proved that $\operatorname{MTL}(\Gamma, \lambda)$ are those models of $\operatorname{MHT}(\Gamma, \lambda)$ satisfying the *excluded middle axiom* schema:

$$\Box_{\ell}(p \lor \neg p) \qquad (\text{for any atom } p \in \mathcal{A}) \tag{5}$$

The semantics of the derived operators in MHT can be easily deduced.

Proposition 1 (Satisfaction)

Let $\mathbf{M} = \langle \mathbf{H}, \mathbf{T} \rangle$ be an HT-trace of length λ over \mathcal{A} . Given the respective definitions of derived operators, we get the following satisfaction conditions:

12. $\mathbf{M}, k \models \blacksquare_n \varphi$ iff for all $j \in [0..\overline{n})$ such that $k - j \in [0..\lambda)$ we have $\mathbf{M}, k - j \models \psi$

13. $\mathbf{M}, k \models \blacklozenge_n \varphi$ iff for some $j \in [0..\overline{n})$ such that $k-j \in [0..\lambda)$ we have $\mathbf{M}, k-j \models \psi$ 14. $\mathbf{M}, k \models \mathbf{I}$ iff k = 0

- 15. $\mathbf{M}, k \models \Box_n \varphi$ iff for all $j \in [0..\overline{n})$ such that $k+j \in [0..\lambda)$ we have $\mathbf{M}, k+j \models \psi$
- 16. $\mathbf{M}, k \models \Diamond_n \varphi$ iff for some $j \in [0, \overline{n})$ such that $k+j \in [0, \lambda)$ we have $\mathbf{M}, k+j \models \psi$
- 17. $\mathbf{M}, k \models \mathbb{F} \text{ iff } k + 1 = \lambda$

18. $\mathbf{M}, k \models \widehat{\bullet} \varphi$ iff k = 0 or $\mathbf{M}, k-1 \models \varphi$

19. $\mathbf{M}, k \models \widehat{\circ}\varphi$ iff $k + 1 = \lambda$ or $\mathbf{M}, k + 1 \models \varphi$

As in the temporal logic of HT (Cabalar et al. 2018), referred to as THT, the operators I and \mathbb{F} exclusively depend on the value of time point k, and are thus independent of **M**. In fact, operator \mathbb{F} allows us to influence the length of models. The inclusion of the axiom $\Diamond_n \mathbb{F}$ for example forces its models to have length $\lambda \leq \overline{n}$ with $\overline{n} \in \mathbb{N}$. On the other hand, the inclusion of the axiom $\neg \Diamond_{\ell} \mathbb{F}$ forces models to be of infinite length. As well, we distinguish MHT on finite and infinite traces, and refer to the respective logics as MHT_f and MHT_{ω} .

Following the definitions of TEL (Cabalar et al. 2018) and DEL (Cabalar et al. 2019), we now introduce non-monotonicity by selecting a particular set of traces that we call *temporal equilibrium models*. First, given an arbitrary set \mathfrak{S} of HT-traces, we define the ones in equilibrium as follows.

Definition 2 (Temporal Equilibrium/Stable models)

Let \mathfrak{S} be some set of HT-traces. A total HT-trace $\langle \mathbf{T}, \mathbf{T} \rangle \in \mathfrak{S}$ is an *equilibrium trace* of \mathfrak{S} iff there is no other $\langle \mathbf{H}, \mathbf{T} \rangle \in \mathfrak{S}$ such that $\mathbf{H} < \mathbf{T}$.

If $\langle \mathbf{T}, \mathbf{T} \rangle$ is such an equilibrium trace, we also say that trace \mathbf{T} is a *stable trace* of \mathfrak{S} . We further talk about *temporal equilibrium* or *temporal stable models* of a theory Γ when $\mathfrak{S} = \mathrm{MHT}(\Gamma)$, respectively.

We write $MEL(\Gamma, \lambda)$ and $MEL(\Gamma)$ to stand for the temporal equilibrium models of $MHT(\Gamma, \lambda)$ and $MHT(\Gamma)$, respectively. Besides, as the ordering relation among traces is only defined for a fixed λ , it is easy to see the following result:

Proposition 2

The set of temporal equilibrium models of Γ can be partitioned by the trace length λ , that is, $\bigcup_{\lambda=0}^{\omega} \text{MEL}(\Gamma, \lambda) = \text{MEL}(\Gamma)$.

Metric Equilibrium Logic (MEL) is the non-monotonic logic induced by temporal equilibrium models of metric theories. We obtain the variants MEL_f and MEL_{ω} by applying the respective restriction to finite or infinite traces, respectively.

Let us illustrate this by using the example of the pedestrian traffic light introduced above. Consider the models of the theory $\Gamma = \{(2), (3), (4)\}$ for length $\lambda = 1$. In this case, we only have time point k = 0, and the metric or temporal aspect is less interesting, since HT-traces amount to pairs $\langle H_0, T_0 \rangle$. Still, this helps to illustrate the difference among the different sets of models defined above. In the example, we abbreviate a set of atoms as a string formed by their initials: for instance gp stands for $\{green, push\}$. Then, we obtain the following sets:

$$\begin{split} \mathrm{MTL}(\Gamma, 1) &= \{ \langle r \rangle, \langle g \rangle, \langle g p \rangle \} \\ \mathrm{MHT}(\Gamma, 1) &= \mathrm{MTL}(\Gamma, 1) \cup \{ \langle \emptyset, g \rangle, \langle \emptyset, g p \rangle, \langle g, g p \rangle \} \\ \mathrm{MEL}(\Gamma, 1) &= \{ \langle r \rangle \} \end{split}$$

As we can see, total models $MTL(\Gamma, 1)$ allow for choosing either *red* or *green* but, if we include *push*, then *green* is mandatory because it is the only way to satisfy $\Diamond_3 \Box_4 green$ with $\lambda = 1$. Note how the unique equilibrium model in $MEL(\Gamma, 1)$ is the only total model $\langle \mathbf{T}, \mathbf{T} \rangle$ with $\mathbf{T} = \langle r \rangle$ in $MTL(\Gamma, 1)$ for which there is no $\langle \mathbf{H}, \mathbf{T} \rangle \in MHT(\Gamma, 1)$ with smaller $\mathbf{H} < \mathbf{T}$. Informally, this is because (3) suffices to justify *red* by default, while the other two total models in $MTL(\Gamma, 1)$, which assume *green* or *push* in T_0 , admit weaker H_0 's where these atoms are not justified. As a result, $MEL(\Gamma, \lambda) = \langle T_i \rangle_{i \in [0..\lambda)}$ with $T_i = \{red\}$ for all $i \in [0..\lambda)$. To illustrate non-monotonicity, suppose we add the formula

$$\circ push$$
 (6)

that ensures the button is pushed in the second state of the trace (i.e. at k = 1) and take $\lambda = 3$. For readability sake, we represent traces (T_0, T_1, T_2) as $T_0 \cdot T_1 \cdot T_2$. For length $\lambda = 3$, formula (4) amounts in MTL to requiring green at k = 2 whenever push holds at any point. Thus, given $\Gamma' = \Gamma \cup \{(6)\}$ and $\lambda = 3$, we get

 $\mathrm{MTL}(\Gamma',3) = \{ \langle \mathbf{T} \rangle = \langle T_0 \cdot T_1 \cdot T_2 \rangle \mid T_0 \in \{r,g,rp,gp\}, T_1 \in \{rp,gp\}, T_2 \in \{g,gp\} \}$

Now, for those $\langle \mathbf{T}, \mathbf{T} \rangle$ with $T_0 \neq r$ and $T_2 = gp$, we always have, among others, a smaller model $\langle \emptyset \cdot T_1 \cdot g, \mathbf{T} \rangle$ in $MHT(\Gamma', 3)$. This means that, in MEL, we conclude by default that

we do not push in other situations $k \neq 1$ and the traffic light is red at the initial state. From the remaining possible total models $\langle r \cdot rp \cdot g \rangle$ and $\langle r \cdot gp \cdot g \rangle$, the latter is not in equilibrium, since $\langle r \cdot p \cdot g, r \cdot gp \cdot g \rangle$ is an model in MHT that reveals that green at k = 0is not justified. As a result MEL($\Gamma', 3$) contains the unique temporal equilibrium model $\langle r \cdot rp \cdot g \rangle$ and the MEL conclusion $\circ^2 r$, we could obtain from Γ alone, is not derived any more once (6) is added to the theory. In this example, we obtain one equilibrium model because $\Box_4 green$ becomes trivially true on traces shorter than $\lambda = 4$. When the trace is long enough ($\lambda \geq 7$), Γ' generates the three expected temporal equilibrium models:

$$\begin{array}{l} \langle r \cdot gp \cdot g \cdot g \cdot g \cdot g \cdot r \cdot r \cdot r & \dots \rangle \\ \langle r \cdot rp \cdot g \cdot g \cdot g \cdot g \cdot g \cdot r \cdot r & \dots \rangle \\ \langle r \cdot rp \cdot r \cdot g \cdot g \cdot g \cdot g \cdot g \cdot r \cdot \dots \rangle \end{array}$$

In the following, we elaborate on the formal characteristics of our approach. At first, we show that a basic property of HT is maintained in MHT:

Proposition 3 (Persistence)

Let $\langle \mathbf{H}, \mathbf{T} \rangle$ be an HT-trace of length λ and φ be a metric formula. Then, for any $k \in [0..\lambda)$,

1. if $\langle \mathbf{H}, \mathbf{T} \rangle$, $k \models \varphi$ then $\langle \mathbf{T}, \mathbf{T} \rangle$, $k \models \varphi$. 2. $\langle \mathbf{H}, \mathbf{T} \rangle$, $k \models \neg \varphi$ iff $\langle \mathbf{T}, \mathbf{T} \rangle$, $k \not\models \varphi$

All MHT tautologies are MTL tautologies but not vice versa (cf. (5) above). However, they coincide for some types of equivalences, as stated below.

Proposition 4

Let φ and ψ be metric formulas without implications (and so, without negations either). Then, $\varphi \equiv \psi$ in MTL iff $\varphi \equiv \psi$ in MHT.

Another useful tool that can save some effort when proving groups of equivalences in MHT is the use of duality properties. A first type of duality has to do with the temporal direction (future or past) of the modal operators. Let \circ / \bullet , $\widehat{\circ} / \widehat{\bullet}$, $\mathbb{U}_n / \mathbf{S}_n$, $\mathbb{R}_n / \mathbf{T}_n$, \Box_n / \blacksquare_n and $\Diamond_n / \blacklozenge_n$ denote all pairs of swapped-time connectives and let $\sigma(\varphi)$ denote the replacement in φ of each connective by its swapped-time version. If we restrict ourselves to finite traces, we get the following result.

Lemma 1

Let **M** be an HT-trace of length λ and φ be a metric formula. Then, there exists an HT-trace **M**' of length λ such that **M**, $k \models \varphi$ iff **M**', $n-k \models \sigma(\varphi)$ for any $k \in [0..\lambda)$.

Theorem 1 (Temporal Duality Theorem)

A metric formula φ is a tautology in MHT_f iff $\sigma(\varphi)$ is a tautology in MHT_f.

For instance, suppose we obtain a proof for

$$\Diamond_n \ p \leftrightarrow p \lor \circ \Diamond_{n-1} \ p \tag{7}$$

Then, we can immediately apply Theorem 1 to guarantee that $\blacklozenge_n p \leftrightarrow p \lor \bullet \blacklozenge_{n-1} p$ is a tautology too. A second kind of duality has to do the analogy between Boolean disjunction and conjunction. Let us define all the pairs of dual connectives as follows: $\land/\lor, \top/\bot$, $\mathbb{U}_n/\mathbb{R}_n$, $\circ/\widehat{\circ}$, \diamondsuit_n/\square_n , $\mathbf{S}_n/\mathbf{T}_n$, $\mathbf{\bullet}/\widehat{\mathbf{\bullet}}$, $\blacklozenge_n/\blacksquare_n$. For a formula φ without implications, we define $\delta(\varphi)$ as the result of replacing each connective by its dual operator. Then, we get the following result.

Proposition 5 (Boolean Duality)

Let φ and ψ be formulas without implication.³ Then, $\varphi \leftrightarrow \psi$ is a tautology in MHT iff $\delta(\varphi) \leftrightarrow \delta(\psi)$ is a tautology in MHT.

Following with our example of equivalence (7), we can now apply Theorem 5 to conclude:

$$\Box_n p \leftrightarrow p \land \bigcirc \Box_{n-1} p$$

Next, we show how metric operators on formulas can be characterized inductively.

Proposition 6

The following formulas are valid in MHT. For any numeral n with $\overline{n} \leq 0$:

$$\varphi \mathbb{U}_n \psi \leftrightarrow \bot \quad \varphi \mathbb{R}_n \psi \leftrightarrow \top \quad \varphi \mathbf{S}_n \psi \leftrightarrow \bot \quad \varphi \mathbf{T}_n \psi \leftrightarrow \top$$

For any numeral n with $\overline{n} > 0$, we have

$$\begin{split} \varphi \mathbb{U}_{n} \psi \leftrightarrow \psi \lor (\varphi \land \bigcirc (\varphi \mathbb{U}_{n-1} \psi)) & \varphi \mathbb{R}_{n} \psi \leftrightarrow \psi \land (\varphi \lor \bigcirc (\varphi \mathbb{R}_{n-1} \psi)) \\ \varphi \mathbf{S}_{n} \psi \leftrightarrow \psi \lor (\varphi \land \blacklozenge (\varphi \mathbf{S}_{n-1} \psi)) & \varphi \mathbf{T}_{n} \psi \leftrightarrow \psi \land (\varphi \lor \bigodot (\varphi \mathbf{G}_{n-1} \psi)) \end{split}$$

The propositions above allow us to unfold metric operators containing numerals. It is easy to see that, for n = 1, the four operators collapse to the formula ψ on their right. For instance, $\Diamond_5 clean \leftrightarrow clean \lor \circ \Diamond_4 clean$ whereas $\Diamond_1 clean \leftrightarrow clean$ and $\Diamond_0 clean \leftrightarrow \bot$.

For metric operators depending on the trace length, the value of which is not necessarily known, Proposition 6 cannot be applied. Instead, we have the following tautologies.

Proposition 7


The following formulas are valid in MHT:

$$\begin{split} \varphi \mathbb{U}_{\ell} \psi \leftrightarrow \psi \lor (\varphi \land \circ (\varphi \mathbb{U}_{\ell} \psi)) & \varphi \mathbb{R}_{\ell} \psi \leftrightarrow \psi \land \left(\varphi \lor \widehat{\circ} (\varphi \mathbb{R}_{\ell} \psi)\right) \\ \varphi \mathbb{S}_{\ell} \psi \leftrightarrow \psi \lor (\varphi \land \bullet (\varphi \mathbb{S}_{\ell} \psi)) & \varphi \mathbb{T}_{\ell} \psi \leftrightarrow \psi \land \left(\varphi \lor \widehat{\circ} (\varphi \mathbb{T}_{\ell} \psi)\right) \end{split}$$

That is, when the limit is the trace length ℓ , the unfolding contains the same operator, it is not altered. As an example, we consider the machine that has to be cleaned eventually before the end of the trace. We then have $\Diamond_{\ell} clean \leftrightarrow clean \lor \circ \Diamond_{\ell} clean$.

Alternatively, metric operators may also be parametrized by intervals⁴ rather than a mere upper bound. In our setting, this is however no restriction since such metric operators can also be expressed, as we show next.

The definition of our interval operators then depends on the type of numeral expression involved. For numeral constants n, m where $\overline{n}, \overline{m} \in [0..\lambda)$, we define:



 $^{^{3}}$ Note that this also means without negation.

⁴ This concept of interval operators should not be confounded with the ones of Allen's interval algebra (Allen 1983).

For intervals spanning to the end (or beginning) of the trace, we have:

As an example of an interval formula, reconsider the machine, discussed in the beginning, and assume that it cannot be cleaned immediately but only within 3 to 5 time steps after usage. This can be expressed as $\Box(use \to \Diamond_{[3;5]} clean)$. In the definitions, note that when $m \ge n$ the interval [m; n) is empty and the operator is always reducible to a truth constant. For instance, $\Diamond_{[5;3]}\varphi$ becomes $\circ^5 \Diamond_{-2}\varphi$ which amounts to $\circ^5 \bot$ or simply \bot . For this reason, we do not consider intervals with ℓ as lower bound, since they are always empty by definition.

Proposition 8 (Satisfaction)

Let $\mathbf{M} = \langle \mathbf{H}, \mathbf{T} \rangle$ be an HT-trace of length λ over \mathcal{A} . Given the respective definitions of derived operators, we get the following satisfaction conditions:

- 20. $\mathbf{M}, k \models \varphi \mathbb{U}_{[m;n)} \psi$ iff for some $j \in [\overline{m}.\overline{n})$ such that $k+j \in [0..\lambda)$ we have $\mathbf{M}, k+j \models \psi$ and $\mathbf{M}, k+i \models \varphi$ for all $i \in [0..j)$
- 21. $\mathbf{M}, k \models \varphi \mathbb{R}_{[m;n)} \psi$ iff for all $j \in [\overline{m}..\overline{n})$ such that $k+j \in [0..\lambda)$, we have $\mathbf{M}, k+j \models \psi$ or $\mathbf{M}, k+i \models \varphi$ for some $i \in [0..j)$
- 22. $\mathbf{M}, k \models \varphi \mathbf{S}_{[m;n)} \psi$ iff for some $j \in [\overline{m}.\overline{n})$ such that $k-j \in [0.\lambda)$ we have $\mathbf{M}, k-j \models \psi$ and $\mathbf{M}, k-i \models \varphi$ for all $i \in [0.j)$
- 23. $\mathbf{M}, k \models \varphi \mathbf{T}_{[m;n)} \psi$ iff for all $j \in [\overline{m}..\overline{n})$ such that $k-j \in [0..\lambda)$, we have $\mathbf{M}, k-j \models \psi$ or $\mathbf{M}, k-i \models \varphi$ for some $i \in [0..j)$
- 24. $\mathbf{M}, k \models \Diamond_{[m:n]} \varphi$ iff for some $j \in [\overline{m}.\overline{n})$ such that $k+j \in [0.\lambda)$ we have $\mathbf{M}, k+j \models \varphi$
- 25. $\mathbf{M}, k \models \Box_{[m:n]} \varphi$ iff for all $j \in [\overline{m}.\overline{n})$ such that $k+j \in [0.\lambda)$, we have $\mathbf{M}, k+j \models \varphi$
- 26. $\mathbf{M}, k \models \oint_{[m;n]} \varphi$ iff for some $j \in [\overline{m}, \overline{n})$ such that $k j \in [0, \lambda)$ we have $\mathbf{M}, k j \models \varphi$
- 27. $\mathbf{M}, k \models \blacksquare_{[m;n]} \varphi$ iff for all $j \in [\overline{m}..\overline{n})$ such that $k-j \in [0..\lambda)$, we have $\mathbf{M}, k-j \models \varphi$

Note that one-step operators can be represented in terms of intervals, since we have

$$\begin{split} & \circ\varphi = \circ \Diamond_1 \varphi = \Diamond_{[1;2)} \varphi & \quad & \bullet\varphi = \bullet \blacklozenge_1 \varphi = \blacklozenge_{[1;2)} \varphi \\ & \hat{\circ}\varphi = \hat{\circ} \Box_1 \varphi = \Box_{[1;2)} \varphi & \quad & \hat{\bullet}\varphi = \hat{\bullet} \blacksquare_1 \varphi = \blacksquare_{[1;2)} \varphi \end{split}$$

Next, we present a three-valued semantics for MHT which turns out to be particularly useful for formal elaborations. In particular, this three-valued interpretation has an important advantage: it allows the interchange of subformulas in a larger formula provided that the interchanged subformulas have the same three-valued interpretations. This characterization relies on temporal three-valued interpretation (Cabalar 2010) and is inspired, in its turn, by the characterization of HT in terms of Gödel's logic G_3 (Gödel 1932). Under this orientation, we deal with three truth values $\{0, 1, 2\}$ standing for: 2 (or proved true) meaning satisfaction "here"; 0 (or assumed false) meaning falsity "there"; and 1 (potentially true) for formulas assumed but not proved to be true. Given an HT-trace $\langle \mathbf{H}, \mathbf{T} \rangle$, we define its associated *truth valuation* as a function $\mathbf{m}(k, \varphi)$ that assigns a truth value in the set $\{0, 1, 2\}$ to a metric formula φ at time point $k \in [0..\lambda)$ as follows. For propositional connectives, $\mathbf{m}(k, \varphi)$ directly corresponds to G_3 , that is, conjunction is the minimum, disjunction the maximum and implication $\boldsymbol{m}(k, \varphi \to \psi)$ is 2 if $\boldsymbol{m}(k, \varphi) \leq \boldsymbol{m}(k, \psi)$, or is $\boldsymbol{m}(k, \psi)$ otherwise. For the rest of cases, we have:

$$\begin{split} \boldsymbol{m}(k, \perp) &\stackrel{\text{def}}{=} & 0 \\ \boldsymbol{m}(k, \top) &\stackrel{\text{def}}{=} & 2 \\ \boldsymbol{m}(k, p) &\stackrel{\text{def}}{=} & \begin{cases} 0 & \text{if } p \notin T_k \\ 1 & \text{if } p \in T_k \setminus H_k & \text{for any atom } p \in \mathcal{A} \\ 2 & \text{if } p \in H_k \end{cases} \\ \boldsymbol{m}(k, \circ \varphi) &\stackrel{\text{def}}{=} & \begin{cases} 0 & \text{if } k+1=\lambda; \\ \boldsymbol{m}(k+1, \varphi) & \text{otherwise} \end{cases} \\ \boldsymbol{m}(k, \widehat{\circ} \varphi) &\stackrel{\text{def}}{=} & \begin{cases} 2 & \text{if } k+1=\lambda; \\ \boldsymbol{m}(k+1, \varphi) & \text{otherwise} \end{cases} \\ \boldsymbol{m}(k+1, \varphi) & \text{otherwise} \end{cases} \\ \boldsymbol{m}(k, \bullet \varphi) &\stackrel{\text{def}}{=} & \begin{cases} 0 & \text{if } k=0; \\ \boldsymbol{m}(k-1, \varphi) & \text{otherwise} \end{cases} \\ \boldsymbol{m}(k, \widehat{\bullet} \varphi) &\stackrel{\text{def}}{=} & \begin{cases} 2 & \text{if } k=0; \\ \boldsymbol{m}(k-1, \varphi) & \text{otherwise} \end{cases} \end{split}$$

For *n* being a numeral constant or the constant ℓ , we have⁵

1 0

$$\begin{split} \boldsymbol{m}(k,\varphi \mathbb{U}_n \psi) & \stackrel{\text{def}}{=} & \max\{\min\{\boldsymbol{m}(k+i,\psi), \boldsymbol{m}(k+j,\varphi) \mid j \in [0..i), \ k+i < \lambda\} \mid 0 \le i < \overline{n}\} \\ \boldsymbol{m}(k,\varphi \mathbb{R}_n \psi) & \stackrel{\text{def}}{=} & \min\{\max\{\boldsymbol{m}(k+i,\psi), \boldsymbol{m}(k+j,\varphi) \mid j \in [0..i), \ k+i < \lambda\} \mid 0 \le i < \overline{n}\} \\ \boldsymbol{m}(k,\varphi \mathbf{S}_n \psi) & \stackrel{\text{def}}{=} & \max\{\min\{\boldsymbol{m}(k-i,\psi), \boldsymbol{m}(k-j,\varphi) \mid j \in [0..i), \ k-i \ge 0\} \mid 0 \le i < \overline{n}\} \\ \boldsymbol{m}(k,\varphi \mathbf{T}_n \psi) & \stackrel{\text{def}}{=} & \min\{\max\{\boldsymbol{m}(k-i,\psi), \boldsymbol{m}(k-j,\varphi) \mid j \in [0..i), \ k-i \ge 0\} \mid 0 \le i < \overline{n}\} \end{split}$$

Proposition 9

Let $\langle \mathbf{H}, \mathbf{T} \rangle$ be a HT-trace of length λ , \boldsymbol{m} its associated valuation and $k \in [0..\lambda)$. Then, for any formula φ , we have

- $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi \text{ iff } \boldsymbol{m}(k, \varphi) = 2 \text{ and }$
- $\langle \mathbf{T}, \mathbf{T} \rangle, k \models \varphi \text{ iff } \boldsymbol{m}(k, \varphi) \neq 0.$

3 Metric and Temporal Equilibrium Logic

In this section, we study the relation between metric and temporal (equilibrium) logics. In fact, temporal formulas constitute a subclass of metric formulas.

The next result guarantees that the original semantics of the temporal operators in THT (Cabalar et al. 2018) is preserved.

⁵ Here, we assume that $\max(\emptyset) = 0$ and $\min(\emptyset) = 2$.

Proposition 10 (Satisfaction)

Let $\mathbf{M} = \langle \mathbf{H}, \mathbf{T} \rangle$ be an HT-trace of length λ over \mathcal{A} . Given the respective definitions of derived operators, we get the following satisfaction conditions:

- 28. $\mathbf{M}, k \models \Diamond \varphi \text{ iff } \mathbf{M}, i \models \varphi \text{ for some } i \in [k..\lambda)$
- 29. $\mathbf{M}, k \models \Box \varphi$ iff $\mathbf{M}, i \models \varphi$ for all $i \in [k..\lambda)$
- 30. $\mathbf{M}, k \models \varphi \mathbb{U} \psi$ iff for some $j \in [k..\lambda)$, we have $\mathbf{M}, j \models \psi$ and $\mathbf{M}, i \models \varphi$ for all $i \in [k..j)$
- 31. $\mathbf{M}, k \models \varphi \mathbb{R} \psi$ iff for all $j \in [k..\lambda)$, we have $\mathbf{M}, j \models \psi$ or $\mathbf{M}, i \models \varphi$ for some $i \in [k..j)$
- 32. $\mathbf{M}, k \models \mathbf{\Phi}\varphi$ iff $\mathbf{M}, i \models \varphi$ for some $i \in [0..k]$
- 33. $\mathbf{M}, k \models \blacksquare \varphi$ iff $\mathbf{M}, i \models \varphi$ for all $i \in [0..k]$
- 34. $\mathbf{M}, k \models \varphi \mathbf{S} \psi$ iff for some $j \in [0..k]$, we have $\mathbf{M}, j \models \psi$ and $\mathbf{M}, i \models \varphi$ for all $i \in (j..k]$
- 35. $\mathbf{M}, k \models \varphi \mathsf{T} \psi$ iff for all $j \in [0..k]$, we have $\mathbf{M}, j \models \psi$ or $\mathbf{M}, i \models \varphi$ for some $i \in (j..k]$

Interestingly, it turns out that metric formulas can also be translated into temporal formulas. This is due to the discrete time domain of MHT and the resulting semantic structure common to MHT and THT. In fact, we provide two alternative translations possessing complementary properties.

Language-preserving translation. Our first translation refrains from extending the original language and is independent of the length of the trace (and so the specific value of ℓ). Although this allows us to search for models of varying length without recompiling a formula, when using temporal ASP solvers such as *telingo* (Cabalar et al. 2019), the translation suffers from an exponential blowup in the worst case.

We define the translation recursively as follows:

$$\tau(a) \stackrel{\text{def}}{=} a \text{ for } a \in \mathcal{A}$$

$$\tau(\oplus \varphi) \stackrel{\text{def}}{=} \oplus \tau(\varphi) \text{ for } \oplus \in \{\neg, \bullet, \circ, \circ, \circ, \circ, \circ\}$$

$$\tau(\varphi \otimes \psi) \stackrel{\text{def}}{=} \tau(\varphi) \otimes \tau(\psi) \text{ for } \otimes \in \{\land, \lor, \rightarrow\}$$

$$\tau(\varphi \otimes_{\ell} \psi) \stackrel{\text{def}}{=} \tau(\varphi) \otimes \tau(\psi) \text{ for } \otimes \in \{\Downarrow, \mathbb{R}, \mathbf{S}, \mathbf{T}\}$$

$$\tau(\varphi \otimes_{1} \psi) \stackrel{\text{def}}{=} \tau(\psi) \text{ for } \otimes \in \{\Downarrow, \mathbb{R}, \mathbf{S}, \mathbf{T}\}$$

$$\tau(\varphi \square_{n} \psi) \stackrel{\text{def}}{=} \tau(\psi \lor (\varphi \land \circ (\varphi \square_{n-1} \psi)))$$

$$\tau(\varphi \mathbf{S}_{n} \psi) \stackrel{\text{def}}{=} \tau(\psi \lor (\varphi \land \bullet (\varphi \mathbf{S}_{n-1} \psi)))$$

$$\tau(\varphi \mathbf{T}_{n} \psi) \stackrel{\text{def}}{=} \tau(\psi \land (\varphi \lor \circ (\varphi \nabla \mathbf{G}_{n-1} \psi)))$$

Translating formula (4) from the traffic light example into a temporal formula yields

$$\tau(\Box(push \to \Diamond_3 \Box_4 green)) = \Box(push \to ((green \land \widehat{\circ} \widehat{\circ} green \land \widehat{\circ} \widehat{\circ} \widehat{\circ} green) \land ((green \land \widehat{\circ} \widehat{\circ} green \land \widehat{\circ} \widehat{\circ} \widehat{\circ} green)) \land (\circ(green \land \widehat{\circ} \widehat{\circ} green \land \widehat{\circ} \widehat{\circ} \widehat{\circ} green)) \land (\circ(green \land \widehat{\circ} \widehat{\circ} green \land \widehat{\circ} \widehat{\circ} \widehat{\circ} green)))))$$

$$(8)$$

This illustrates the benefit of metric operators. While we are also able to express the

10

same formula with temporal operators, it is much more concise and more readable with metric operators.

Proposition 11 The translation $\tau(\varphi)$ terminates for any metric formula φ .

Proposition 12

For any HT-trace **M** and any time point $k \in [0..\lambda)$, we have **M**, $k \models \varphi$ in MHT iff **M**, $k \models \tau(\varphi)$ in THT.

Corollary 1

For any metric formula φ over \mathcal{A} , there exists a temporal formula ψ over \mathcal{A} such that an HT-trace **M** is a model for ψ in THT iff **M** is a model for φ in MHT.

Language-extending translation. For a complement, we provide an alternative translation using an extended alphabet. This translation has the advantage of avoiding applications of distributivity, a source of an exponential increase in size.

To this end, we adapt the notion of closure from (Cabalar et al. 2020) to provide a translation of metric into temporal formulas. The original definition of closure is due to (Fischer and Ladner 1979).

Definition 3 (Closure)

The closure $cl(\gamma)$ of a metric formula γ is the subset minimal set of formulas satisfying the inductive conditions:

1. $\gamma \in \operatorname{cl}(\gamma)$ 2. $(\varphi \otimes \psi) \in \operatorname{cl}(\gamma)$ implies $\varphi \in \operatorname{cl}(\gamma)$ and $\psi \in \operatorname{cl}(\gamma)$ for $\otimes \in \{\wedge, \vee, \rightarrow, \bigcup_n, \mathbb{R}_n, \mathbf{S}_n, \mathbf{T}_n\}$ with $n \in \mathbb{N} \cup \boldsymbol{\ell}$ 3. If $\otimes \psi \in \operatorname{cl}(\gamma)$ then $\psi \in \operatorname{cl}(\gamma)$ for $\otimes \in \{\circ, \circ, \bullet, \bullet\}$ 4. If $\varphi \bigcup_n \psi \in \operatorname{cl}(\gamma)$ and n > 1 then $\circ (\varphi \bigcup_{n-1} \psi) \in \operatorname{cl}(\gamma)$ 5. If $\varphi \mathbb{R}_n \psi \in \operatorname{cl}(\gamma)$ and n > 1 then $\circ (\varphi \mathbb{R}_{n-1} \psi) \in \operatorname{cl}(\gamma)$ 6. If $\varphi \mathbb{S}_n \psi \in \operatorname{cl}(\gamma)$ and n > 1 then $\bullet (\varphi \mathbb{S}_{n-1} \psi) \in \operatorname{cl}(\gamma)$ 7. If $\varphi \mathbb{T}_n \psi \in \operatorname{cl}(\gamma)$ and n > 1 then $\bullet (\varphi \mathbb{S}_{n-1} \psi) \in \operatorname{cl}(\gamma)$ 8. If $\varphi \bigcup_{\ell} \psi \in \operatorname{cl}(\gamma)$ then $\circ (\varphi \bigcup_{\ell} \psi) \in \operatorname{cl}(\gamma)$ 9. If $\varphi \mathbb{R}_{\ell} \psi \in \operatorname{cl}(\gamma)$ then $\circ (\varphi \mathbb{R}_{\ell} \psi) \in \operatorname{cl}(\gamma)$ 10. If $\varphi \mathbb{S}_{\ell} \psi \in \operatorname{cl}(\gamma)$ then $\bullet (\varphi \mathbb{S}_{\ell} \psi) \in \operatorname{cl}(\gamma)$ 11. If $\varphi \mathbb{T}_{\ell} \psi \in \operatorname{cl}(\gamma)$ then $\widehat{\bullet} (\varphi \mathbb{T}_{\ell} \psi) \in \operatorname{cl}(\gamma)$

Any set satisfying these conditions is called *closed*.

Proposition 13

For any metric formula φ , $cl(\varphi)$ is finite. Moreover, the total size of all the formulas in $cl(\varphi)$, $|cl(\varphi)|$ is bound by $|cl(\varphi)| \leq 2k_{\varphi}|\varphi|$, where $k_{\varphi} = \max\{1, n_{\varphi}\}$ and n_{φ} is the maximum (metric) integer subindex occurring in φ .

Thus, given a metric formula φ over alphabet \mathcal{A} at hand, we define the extended alphabet $\mathcal{A}^{\varphi} \stackrel{def}{=} \mathcal{A} \cup \{\mathbf{L}_{\mu} \mid \mu \in \mathrm{cl}(\varphi)\}$. For convenience, we simply use $\mathbf{L}_{\varphi} \stackrel{def}{=} \varphi$ if φ is \top, \bot or an atom $a \in \mathcal{A}$.

As happened with the normal form reduction for TEL_f in (Cabalar et al. 2018), the

translation is done in two phases: we first obtain a temporal theory containing double implications, and then we unfold them into temporal rules. We start by defining the temporal theory $v(\varphi)$ that introduces new labels \mathbf{L}_{μ} for each formula $\mu \in cl(\varphi)$. This theory contains the formula \mathbf{L}_{φ} and, per each label \mathbf{L}_{μ} , a set of formulas $df(\mu)$ fixing the label's truth value. Formally, we define that

$$v(\varphi) = \{\mathbf{L}_{\varphi}\} \cup \{df(\mu) \mid \mu \in \mathrm{cl}(\varphi)\} \qquad \text{and} \qquad v(\Gamma) = \{v(\varphi) \mid \varphi \in \Gamma\}.$$

Table 1 shows the definitions $df(\mu)$ for each μ in the closure $cl(\varphi)$ depending on the outer modality in the formula.

r	1
$\mu\in \mathrm{cl}(\varphi)$	$df(\mu)$
Οα	$\widehat{O}\Box(\mathbf{\bullet}\mathbf{L}_{\mu}\leftrightarrow\mathbf{L}_{\alpha})\qquad \Box(\mathbb{F}\rightarrow\neg\mathbf{L}_{\mu})$
$\bullet \alpha$	$\widehat{O}\Box(\mathbf{L}_{\mu}\leftrightarrow ullet \mathbf{L}_{lpha}) ~~ abla \mathbf{L}_{\mu}$
Ôα	$\widehat{O}\square(\mathbf{ullet}\mathbf{L}_\mu\leftrightarrow\mathbf{L}_lpha)$ $\square(\mathbb{F} ightarrow\mathbf{L}_\mu)$
$\widehat{\bullet}\alpha$	$\widehat{O}\Box(\mathbf{L}_{\mu}\leftrightarrow ullet \mathbf{L}_{lpha}) \qquad \mathbf{L}_{\mu}$
$\alpha \mathbb{U}_1 \beta$	$\Box(\mathbf{L}_{\mu}\leftrightarrow\mathbf{L}_{eta})$
$\alpha \mathbb{R}_1 \beta$	$\Box(\mathbf{L}_{\mu}\leftrightarrow\mathbf{L}_{eta})$
$\alpha \mathbf{S}_1 \beta$	$\Box(\mathbf{L}_{\mu}\leftrightarrow\mathbf{L}_{eta})$
$\alpha \mathbf{T}_1 \beta$	$\Box(\mathbf{L}_{\mu}\leftrightarrow\mathbf{L}_{eta})$
$\alpha U_n \beta$	$\Box (\mathbf{L}_{\mu} \leftrightarrow \mathbf{L}_{\beta} \lor (\mathbf{L}_{\alpha} \land \mathbf{L}_{\alpha'})) \qquad \text{with } \alpha' = O (\alpha \mathbb{U}_{n-1}\beta)$
$\alpha \mathbb{R}_n \beta$	$\Box (\mathbf{L}_{\mu} \leftrightarrow \mathbf{L}_{\beta} \land (\mathbf{L}_{\alpha} \lor \mathbf{L}_{\alpha'})) \qquad \text{with } \alpha' = \widehat{O} (\alpha \mathbb{R}_{n-1}\beta)$
$\alpha \mathbf{S}_n \beta$	$\Box (\mathbf{L}_{\mu} \leftrightarrow \mathbf{L}_{\beta} \lor (\mathbf{L}_{\alpha} \land \mathbf{L}_{\alpha'})) \qquad \text{with } \alpha' = \mathbf{\bullet} (\alpha \mathbf{S}_{n-1}\beta)$
$\alpha \mathbf{T}_n \beta$	$\Box (\mathbf{L}_{\mu} \leftrightarrow \mathbf{L}_{\beta} \wedge (\mathbf{L}_{\alpha} \vee \mathbf{L}_{\alpha'})) \text{with } \alpha' = \widehat{\bullet} (\alpha T_{n-1}\beta)$
$\alpha \mathbb{U}_{\ell} \beta$	$\Box (\mathbf{L}_{\mu} \leftrightarrow \mathbf{L}_{\beta} \lor (\mathbf{L}_{\alpha} \land \mathbf{L}_{\alpha'})) \qquad \text{with } \alpha' = O (\alpha \mathbb{U}_{\ell} \beta)$
$\alpha \mathbb{R}_{\ell} \beta$	$\Box (\mathbf{L}_{\mu} \leftrightarrow \mathbf{L}_{\beta} \land (\mathbf{L}_{\alpha} \lor \mathbf{L}_{\alpha'})) \qquad \text{with } \alpha' = \widehat{O} (\alpha \mathbb{R}_{\boldsymbol{\ell}} \beta)$
$\alpha S_{\ell} \beta$	$\Box (\mathbf{L}_{\mu} \leftrightarrow \mathbf{L}_{\beta} \lor (\mathbf{L}_{\alpha} \land \mathbf{L}_{\alpha'})) \qquad \text{with } \alpha' = \mathbf{\bullet} (\alpha \mathbf{S}_{\boldsymbol{\ell}} \beta)$
$\alpha \mathbf{T}_{\boldsymbol{\ell}} \beta$	$\Box (\mathbf{L}_{\mu} \leftrightarrow \mathbf{L}_{\beta} \wedge (\mathbf{L}_{\alpha} \vee \mathbf{L}_{\alpha'})) \qquad \text{with } \alpha' = \widehat{\bullet} (\alpha T_{\boldsymbol{\ell}} \beta)$

Table 1. Translation of metric modal operators

As we have seen, in the general case, formulas in $df(\mu)$ are not temporal rules, since they sometimes contain double implications. However, they all have the forms φ , $\Box \varphi$, $\widehat{\circ} \Box \varphi$ or $\Box (\mathbb{F} \to \varphi)$, for some inner propositional formula φ formed with temporal literals.

Given an HT-trace $\langle \mathbf{H}, \mathbf{T} \rangle = \langle H_i, T_i \rangle_{i \in [0..\lambda)}$, we define its restriction to alphabet \mathcal{A} as $\langle \mathbf{H}, \mathbf{T} \rangle |_{\mathcal{A}} \stackrel{def}{=} \langle H_i \cap \mathcal{A}, T_i \cap \mathcal{A} \rangle_{i \in [0..\lambda)}$. Similarly, for any set \mathfrak{S} of HT-traces, we write $\mathfrak{S}|_{\mathcal{A}}$ to stand for $\{ \langle \mathbf{H}, \mathbf{T} \rangle |_{\mathcal{A}} | \langle \mathbf{H}, \mathbf{T} \rangle \in \mathfrak{S} \}$.

The following lemma shows that \mathbf{L}_{μ} and μ are equivalent:

Proposition 14

Let γ be a metric formula over \mathcal{A} and let $\langle \mathbf{H}, \mathbf{T} \rangle$ be a model in MHT_f of $v(\gamma)$ being associated with the three-valuation \boldsymbol{m} .

Then, for any $\mu \in cl(\gamma)$ and any $k \in [0..\lambda)$, we have $\boldsymbol{m}(k, \mathbf{L}_{\mu}) = \boldsymbol{m}(k, \mu)$.

Theorem 2

For any metric formula φ and any length λ , we have

$$\mathrm{MHT}_f(\varphi, \lambda) = \mathrm{MHT}_f(\upsilon(\varphi), \lambda)|_{\mathcal{A}} = \mathrm{THT}_f(\upsilon(\varphi), \lambda)|_{\mathcal{A}}.$$

Corollary 2

For any temporal formula φ and any length λ , we have

$$THT(\varphi, \lambda) = MHT(\varphi, \lambda)$$

For example, the formula $\Box(\neg green \rightarrow red)$ in (3) has the same models whether it is interpreted as a temporal or a metric formula.

Corollary 3

Let φ be a metric formula over \mathcal{A} . Then, translation $v(\varphi)$ is strongly faithful, that is:

$$\operatorname{MEL}_f(\varphi \wedge \varphi') = \operatorname{MEL}_f(\upsilon(\varphi) \wedge \varphi')|_{\mathcal{A}}.$$

for any arbitrary metric formula φ' over \mathcal{A} .

The translation of formula (4) from the traffic light example yields the resulting set of rules in Table 2 computed from cl(4). Although in this case, cl(4) is larger than the

\mathbf{L}_{μ}	$\mu \in cl(4)$	$df(\mu)$
\mathbf{L}_1	$\Box (push \to \Diamond_3 \Box_4 green)$	$\Box \left(\mathbf{L}_{1} \leftrightarrow \mathbf{L}_{3} \wedge \mathbf{L}_{2} ight)$
\mathbf{L}_2	$\widehat{O}\Box\left(push \to \Diamond_3\Box_4 green\right)$	$\widehat{O}\Box\left(\mathbf{\bullet}\mathbf{L}_{2}\leftrightarrow\mathbf{L}_{1}\right)\qquad\Box\left(\mathbb{F}\rightarrow\mathbf{L}_{2}\right)$
\mathbf{L}_3	$push \rightarrow \Diamond_3 \Box_4 green$	$\Box \left(\mathbf{L}_{3} \leftrightarrow (push \rightarrow \mathbf{L}_{4}) \right)$
\mathbf{L}_4	$\Diamond_3 \Box_4 green$	$\Box \left(\mathbf{L}_{4} \leftrightarrow \mathbf{L}_{9} \lor \mathbf{L}_{5} ight)$
\mathbf{L}_5	$O\Diamond_2\Box_4$ green	$\widehat{O}\Box\left(\mathbf{\bullet}\mathbf{L}_{5}\leftrightarrow\mathbf{L}_{6}\right)\qquad\Box\left(\mathbb{F}\rightarrow\neg\mathbf{L}_{5}\right)$
\mathbf{L}_{6}	$\Diamond_2 \Box_4 green$	$\Box \left(\mathbf{L}_{6} \leftrightarrow \mathbf{L}_{9} \lor \mathbf{L}_{7} \right)$
\mathbf{L}_7	$O\Diamond_1\Box_4$ green	$\widehat{O}\Box\left(\mathbf{\bullet}\mathbf{L}_{7}\leftrightarrow\mathbf{L}_{8}\right)\qquad\Box\left(\mathbb{F}\rightarrow\neg\mathbf{L}_{7}\right)$
\mathbf{L}_8	$\Diamond_1 \Box_4 green$	$\Box \left(\mathbf{L}_{8} \leftrightarrow \mathbf{L}_{9} ight)$
\mathbf{L}_9	$\Box_4 green$	$\Box \left(\mathbf{L}_{9} \leftrightarrow green \wedge \mathbf{L}_{10} \right)$
\mathbf{L}_{10}	$\widehat{O}\square_3$ green	$\widehat{O}\Box\left(\mathbf{\bullet}\mathbf{L}_{10}\leftrightarrow\mathbf{L}_{11}\right)\qquad\Box\left(\mathbb{F}\rightarrow\mathbf{L}_{10}\right)$
\mathbf{L}_{11}	$\Box_3 green$	$\Box \left(\mathbf{L}_{11} \leftrightarrow green \wedge \mathbf{L}_{12} \right)$
\mathbf{L}_{12}	$\widehat{O}\square_2 green$	$\widehat{O}\Box\left(\mathbf{\bullet}\mathbf{L}_{12}\leftrightarrow\mathbf{L}_{13}\right)\qquad\Box\left(\mathbb{F}\rightarrow\mathbf{L}_{12}\right)$
\mathbf{L}_{13}	$\Box_2 green$	$\Box \left(\mathbf{L}_{13} \leftrightarrow green \wedge \mathbf{L}_{14} \right)$
\mathbf{L}_{14}	$\widehat{O}\Box_1 green$	$\widehat{O}\Box\left(\mathbf{\bullet}\mathbf{L}_{14}\leftrightarrow\mathbf{L}_{15}\right)\qquad\Box\left(\mathbb{F}\rightarrow\mathbf{L}_{14}\right)$
\mathbf{L}_{15}	$\Box_1 green$	$\Box \left(\mathbf{L}_{15} \leftrightarrow green \right)$

Table 2. Translating formula (4) from the traffic light example.

language-preserving translation $\tau(4) = (8)$, note that the latter is not in the form of a logic program yet, while double implications in Table 2 can be linearly reduced to a logic program. In general, reducing $\tau(\gamma)$ to a (language-preserving) logic program induces an exponential size increase due to the application of distributivity laws. For instance, the formula $\tau(p \bigcup_n q)$ amounts to a combination of conjunctions and disjunctions that, to become a logic program, must be previously reduced to conjunctive normal form. On the other hand, $cl(\gamma)$ preserves a polynomial size, although at the cost of introducing auxiliary atoms, one per formula in $cl(\gamma)$.

4 Discussion

We have defined and elaborated upon a metric temporal extension of the logic of Hereand-There in order to lay the theoretical foundations of metric Answer Set Programming. The resulting logics, MHT and its non-monotonic extension MEL, have a point-based semantics based on discrete linear time. The choice of such a simple time domain was motivated by the desire to base the approach on the same semantic structures as used in previous extensions of HT and ASP with constructs from Linear Temporal and Dynamic Logic, namely, linear HT-traces. As a result, we were able to inter-translate our approach and the temporal logics of THT and TEL. This is of practical relevance since it allows us to use temporal ASP solvers such as *telingo* (Cabalar et al. 2019) for implementation.

There exist other approaches that introduce metric temporal operators in logic programming. For instance, a first metric extension for Horn Prolog was presented back in (Brzoska 1995). This approach is more limited than MEL for several reasons: it does not consider default negation, metric operators can only be used under a restricted syntax and only for unary temporal operators, and the interpretation of programs relies on resolution rather than on a purely model-theoretic description. Other more recent related approaches introduce temporal metrics for stream reasoning for ASP (Beck et al. 2015; Beck et al. 2016) and DATALOG (Brandt et al. 2018; Walega et al. 2019). The former is not only closely related due to its ASP-based approach but also because it can be characterized in terms of EL (Beck et al. 2016), although extra conditions are needed to guarantee the persistence property of HT. While the DATALOG-based approach lacks the rich language of ASP, they rely on more expressive metric operators. For instance, (Brandt et al. 2018) deal with intervals over $\mathbb{Q} \cup \{-\infty, +\infty\}$. It will be interesting to investigate the formal connection to such stream-oriented approaches and metric temporal logic programming formalisms in general.

Another aspect to explore has to do with the fact that traditional metric logics rely on continuous time domains (cf. (Alur and Henzinger 1992)), which often brings about undecidability. Although we do not strive for such expressiveness, it will be interesting future work to generalize our approach to more fine-grained time domains using integer or rational numbers and to explore implementations with hybrid ASP systems. However, the presented approach should already furnish a host system for action languages with durative actions (Son et al. 2004) or even traditional event calculus (Kowalski and Sergot 1986), although this remains to be worked out.

References

- AGUADO, F., CABALAR, P., DIÉGUEZ, M., PÉREZ, G., AND VIDAL, C. 2013. Temporal equilibrium logic: a survey. *Journal of Applied Non-Classical Logics 23*, 1-2, 2–24.
- ALLEN, J. 1983. Maintaining knowledge about temporal intervals. Communications of the ACM 26, 11, 832–843.
- ALUR, R. AND HENZINGER, T. 1992. Logics and models of real time: A survey. In *Real-Time: Theory in Practice*. Springer, 74–106.
- BALDUCCINI, M., LIERLER, Y., AND WOLTRAN, S., Eds. 2019. Proceedings of the Fifteenth International Conference on Logic Programming and Nonmonotonic Reasoning (LPNMR'19). Springer.

BECK, H., DAO-TRAN, M., AND EITER, T. 2016. Equivalent stream reasoning programs.

In Proceedings of the Twenty-fifth International Joint Conference on Artificial Intelligence (IJCAI'16), R. Kambhampati, Ed. IJCAI/AAAI Press, 929–935.

- BECK, H., DAO-TRAN, M., EITER, T., AND FINK, M. 2015. LARS: A logic-based framework for analyzing reasoning over streams. In *Proceedings of the Twenty-Ninth National Conference on Artificial Intelligence (AAAI'15)*, B. Bonet and S. Koenig, Eds. AAAI Press, 1431–1438.
- BOSSER, A., CABALAR, P., DIÉGUEZ, M., AND SCHAUB, T. 2018. Introducing temporal stable models for linear dynamic logic. In *Proceedings of the Sixteenth International Conference on Principles of Knowledge Representation and Reasoning (KR'18)*, M. Thielscher, F. Toni, and F. Wolter, Eds. AAAI Press, 12–21.
- BRANDT, S., KALAYCI, E., RYZHIKOV, V., XIAO, G., AND ZAKHARYASCHEV, M. 2018. Querying log data with metric temporal logic. *Journal of Artificial Intelligence Research* 62, 829–877.
- BRZOSKA, C. 1995. Temporal logic programming with metric and past operators. In Proceedings of the Workshop on Executable Modal and Temporal Logics, M. Fisher and R. Owens, Eds. Springer, 21–39.
- CABALAR, P. 2010. A normal form for linear temporal equilibrium logic. In Proceedings of the Twelfth European Conference on Logics in Artificial Intelligence (JELIA'10), T. Janhunen and I. Niemelä, Eds. Springer, 64–76.
- CABALAR, P., DIÉGUEZ, M., LAFERRIERE, F., AND SCHAUB, T. 2020. Implementing dynamic answer set programming over finite traces. In *Proceedings of the Twenty-fourth European Conference on Artificial Intelligence (ECAI'20)*, G. De Giacomo, Ed. IOS Press. To appear.
- CABALAR, P., DIÉGUEZ, M., AND SCHAUB, T. 2019. Towards dynamic answer set programming over finite traces. See Balduccini et al. (2019), 148–162.
- CABALAR, P., KAMINSKI, R., MORKISCH, P., AND SCHAUB, T. 2019. telingo = ASP + time. See Balduccini et al. (2019), 256–269.
- CABALAR, P., KAMINSKI, R., SCHAUB, T., AND SCHUHMANN, A. 2018. Temporal answer set programming on finite traces. *Theory and Practice of Logic Programming* 18, 3-4, 406–420.
- FISCHER, M. AND LADNER, R. 1979. Propositional dynamic logic of regular programs. Journal of Computer and System Sciences 18, 2, 194–211.
- GEBSER, M., KAMINSKI, R., KAUFMANN, B., OSTROWSKI, M., SCHAUB, T., AND WANKO, P. 2016. Theory solving made easy with clingo 5. In *Technical Communications of the Thirty*second International Conference on Logic Programming (ICLP'16), M. Carro and A. King, Eds. OpenAccess Series in Informatics (OASIcs), 2:1–2:15.
- GÖDEL, K. 1932. Zum intuitionistischen Aussagenkalkül. Anzeiger der Akademie der Wissenschaften in Wien, 65–66.
- HAREL, D., TIURYN, J., AND KOZEN, D. 2000. Dynamic Logic. MIT Press.
- HEYTING, A. 1930. Die formalen Regeln der intuitionistischen Logik. In Sitzungsberichte der Preussischen Akademie der Wissenschaften. Deutsche Akademie der Wissenschaften zu Berlin, 42–56. Reprint in Logik-Texte: Kommentierte Auswahl zur Geschichte der Modernen Logik, Akademie-Verlag, 1986.
- KOWALSKI, R. AND SERGOT, M. 1986. A logic-based calculus of events. New Generation Computing 4, 1, 67–95.
- LIFSCHITZ, V. 1999. Answer set planning. In Proceedings of the International Conference on Logic Programming (ICLP'99), D. de Schreye, Ed. MIT Press, 23–37.
- OUAKNINE, J. AND WORRELL, J. 2005. On the decidability of metric temporal logic. In Proceedings of the Twentieth Annual Symposium on Logic in Computer Science (LICS'10). IEEE Computer Society Press, 188–197.
- PEARCE, D. 1997. A new logical characterisation of stable models and answer sets. In Proceedings of the Sixth International Workshop on Non-Monotonic Extensions of Logic Programming (NMELP'96), J. Dix, L. Pereira, and T. Przymusinski, Eds. Springer, 57–70.

- PNUELI, A. 1977. The temporal logic of programs. In *Proceedings of the Eight-teenth Symposium* on Foundations of Computer Science (FOCS'77). IEEE Computer Society Press, 46–57.
- SON, T., BARAL, C., AND TUAN, L. 2004. Adding time and intervals to procedural and hierarchical control specifications. In *Proceedings of the Nineteenth National Conference on Artificial Intelligence (AAAI'04)*, D. McGuinness and G. Ferguson, Eds. AAAI Press, 92–97.
- WALEGA, P., KAMINSKI, M., AND CUENCA GRAU, B. 2019. Reasoning over streaming data in metric temporal datalog. In Proceedings of the Thirty-third National Conference on Artificial Intelligence (AAAI'19), P. Van Hentenryck and Z. Zhou, Eds. AAAI Press, 3092–3099.

Appendix A Proofs

Proof of Proposition 3. The first item is proved by structural induction. We consider the different cases below:

- Case $\varphi \mathbb{U}_n \psi$: if $\langle \mathbf{H}, \mathbf{T} \rangle$, $k \models \varphi \mathbb{U}_n \psi$ then there exists $0 \leq i < \overline{n}$ such that $k + i < \lambda$, $\langle \mathbf{H}, \mathbf{T} \rangle$, $k + i \models \psi$ and for all $0 \leq j < i$, $\langle \mathbf{H}, \mathbf{T} \rangle$, $k + j \models \varphi$. By induction hypothesis, $\langle \mathbf{T}, \mathbf{T} \rangle$, $k + i \models \psi$ and for all $0 \leq j < i$, $\langle \mathbf{T}, \mathbf{T} \rangle$, $k + j \models \varphi$. Therefore, $\langle \mathbf{H}, \mathbf{T} \rangle$, $k \models \varphi \mathbb{U}_n \psi$.
- Case $\varphi \mathbb{R}_n \psi$: assume by contradiction that $\langle \mathbf{T}, \mathbf{T} \rangle, k \not\models \varphi \mathbb{R}_n \psi$. This means that there exists $0 \leq i < \overline{n}$ such that $k + i < \lambda$, $\langle \mathbf{T}, \mathbf{T} \rangle, k + i \not\models \psi$ and for all $0 \leq j < i$, $j < i, \langle \mathbf{T}, \mathbf{T} \rangle, k + j \not\models \varphi$. By induction, $\langle \mathbf{H}, \mathbf{T} \rangle, k + i \not\models \psi$ and for all $0 \leq j < i$, $\langle \mathbf{H}, \mathbf{T} \rangle, k + j \not\models \varphi$. Therefore, $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \mathbb{R}_n \psi$: a contradiction.
- Case $\varphi \mathbf{S}_n \psi$: if $\langle \mathbf{H}, \mathbf{T} \rangle$, $k \models \varphi \mathbf{S}_n \psi$ then there exists $0 \leq i < \overline{n}$ such that $k i \geq 0$, $\langle \mathbf{H}, \mathbf{T} \rangle$, $k - i \models \psi$ and for all $0 \leq j < i$, $\langle \mathbf{H}, \mathbf{T} \rangle$, $k - j \models \varphi$. By the induction hypothesis, $\langle \mathbf{T}, \mathbf{T} \rangle$, $k - i \models \psi$ and for all $0 \leq j < i$, $\langle \mathbf{T}, \mathbf{T} \rangle$, $k - j \models \varphi$. Therefore, $\langle \mathbf{T}, \mathbf{T} \rangle$, $k \models \varphi \mathbf{S}_n \psi$
- Case $\varphi \mathbf{T}_n \psi$: assume by contradiction that $\langle \mathbf{T}, \mathbf{T} \rangle, k \not\models \varphi \mathbf{T}_n \psi$. This means that there exists $0 \leq i < \overline{n}$ such that $k i \geq 0$, $\langle \mathbf{T}, \mathbf{T} \rangle, k i \not\models \psi$ and for all $0 \leq j < i$, $\langle \mathbf{T}, \mathbf{T} \rangle, k j \not\models \varphi$. By induction hypothesis, $\langle \mathbf{H}, \mathbf{T} \rangle, k i \not\models \psi$ and for all $0 \leq j < i$, $\langle \mathbf{H}, \mathbf{T} \rangle, k j \not\models \varphi$, so $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \mathbf{T}_n \psi$: a contradiction.
- Case $\varphi \bigcup_{\ell} \psi$: if $\langle \mathbf{H}, \mathbf{T} \rangle$, $k \models \varphi \bigcup_{\ell} \psi$ then there exists $0 \le i < \lambda$ such that $k + i < \lambda$, $\langle \mathbf{H}, \mathbf{T} \rangle$, $k + i \models \psi$ and for all $0 \le j < i$, $\langle \mathbf{H}, \mathbf{T} \rangle$, $k + j \models \varphi$. By induction hypothesis, $\langle \mathbf{T}, \mathbf{T} \rangle$, $k + i \models \psi$ and for all $0 \le j < i$, $\langle \mathbf{T}, \mathbf{T} \rangle$, $k + j \models \varphi$. Therefore, $\langle \mathbf{H}, \mathbf{T} \rangle$, $k \models \varphi \bigcup_{\ell} \psi$.
- Case $\varphi \mathbb{R}_{\ell} \psi$: assume by contradiction that $\langle \mathbf{T}, \mathbf{T} \rangle, k \not\models \varphi \mathbb{R}_{\ell} \psi$. This means that there exists $0 \leq i < \lambda$ such that $k + i < \lambda, \langle \mathbf{T}, \mathbf{T} \rangle, k + i \not\models \psi$ and for all $0 \leq j < i, \langle \mathbf{T}, \mathbf{T} \rangle, k + j \not\models \varphi$. By induction, $\langle \mathbf{H}, \mathbf{T} \rangle, k + i \not\models \psi$ and for all $0 \leq j < i, \langle \mathbf{H}, \mathbf{T} \rangle, k + j \not\models \varphi$. Therefore, $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \mathbb{R}_{\ell} \psi$: a contradiction.
- Case $\varphi \mathbf{S}_{\boldsymbol{\ell}} \psi$: if $\langle \mathbf{H}, \mathbf{T} \rangle$, $k \models \varphi \mathbf{S}_{\boldsymbol{\ell}} \psi$ then there exists $0 \leq i < \lambda$ such that $k i \geq 0$, $\langle \mathbf{H}, \mathbf{T} \rangle$, $k - i \models \psi$ and for all $0 \leq j < i$, $\langle \mathbf{H}, \mathbf{T} \rangle$, $k - j \models \varphi$. By the induction hypothesis, $\langle \mathbf{T}, \mathbf{T} \rangle$, $k - i \models \psi$ and for all $0 \leq j < i$, $\langle \mathbf{T}, \mathbf{T} \rangle$, $k - j \models \varphi$. Therefore, $\langle \mathbf{T}, \mathbf{T} \rangle$, $k \models \varphi \mathbf{S}_{\boldsymbol{\ell}} \psi$
- Case $\varphi \mathbf{T}_{\boldsymbol{\ell}} \psi$: assume by contradiction that $\langle \mathbf{T}, \mathbf{T} \rangle, k \not\models \varphi \mathbf{T}_{\boldsymbol{\ell}} \psi$. This means that there exists $0 \leq i < \lambda$ such that $k i \geq 0$, $\langle \mathbf{T}, \mathbf{T} \rangle, k i \not\models \psi$ and for all $0 \leq j < i$, $\langle \mathbf{T}, \mathbf{T} \rangle, k j \not\models \varphi$. By induction hypothesis, $\langle \mathbf{H}, \mathbf{T} \rangle, k i \not\models \psi$ and for all $0 \leq j < i$, $\langle \mathbf{H}, \mathbf{T} \rangle, k j \not\models \varphi$, so $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \mathbf{T}_{\boldsymbol{\ell}} \psi$: a contradiction.

For the second item, assume by contradiction that $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \neg \varphi$ but $\langle \mathbf{T}, \mathbf{T} \rangle, k \models \varphi$. By persistency, $\langle \mathbf{T}, \mathbf{T} \rangle, k \models \neg \varphi$ and, therefore, $\langle \mathbf{T}, \mathbf{T} \rangle, k \models \bot$: a contradiction. Conversely, Assume by contradiction that $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \neg \varphi$. Therefore, either $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi$ or $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi$. In any case, by persistence, we conclude that $\langle \mathbf{T}, \mathbf{T} \rangle, k \not\models \varphi$: a contradiction.

QED

Proof of Proposition 6.

For $\varphi \mathbb{U}_1 \psi \leftrightarrow \psi$ we reason as follows: from left to right, if $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi \mathbb{U}_1 \psi$ then $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \psi$ by definition. Conversely, if $\langle \mathbf{H}, \mathbf{T} \rangle, k + 0 \models \psi$ then $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi \mathbb{U}_1 \psi$ by definition.

The proofs for $\varphi \mathbb{R}_1 \psi \leftrightarrow \psi$, $\varphi \mathbf{S}_1 \psi \leftrightarrow \psi$ and $\varphi \mathbf{T}_1 \psi \leftrightarrow \psi$ are done in a similar way. From now on we will consider n with $\overline{n} > 1$.

• For $\varphi \mathbb{U}_n \psi \leftrightarrow \psi \lor (\varphi \land \circ \varphi \mathbb{U}_{n-1} \psi)$, let us consider from left to right that $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi \mathbb{U}_n \psi$. This means that there exists $0 \leq i < \overline{n}$ such that $k+i < \lambda, \langle \mathbf{H}, \mathbf{T} \rangle, k+i \models \psi$ and for all $0 \leq j < i, \langle \mathbf{H}, \mathbf{T} \rangle, k+j \models \varphi$. If i = 0 then we can easily conclude $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \psi$. If i > 0 then $\langle \mathbf{H}, \mathbf{T} \rangle, k+1 \models \varphi \mathbb{U}_{n-1} \psi$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi$. As a consequence, $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \psi \lor (\varphi \land \circ (\varphi \mathbb{U}_{n-1} \psi))$.

For the converse direction, assume that $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \psi \lor (\varphi \land \circ (\varphi \bigcup_{n-1} \psi))$. If $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \psi$ then $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi \bigcup_n \psi$ by definition (just take i = 0). If $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi \land \circ (\varphi \bigcup_{n-1} \psi)$ then $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi, k+1 < \lambda$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k+1 \models \varphi \bigcup_{n-1} \psi$. This means that there exists $0 \le i < \overline{n-1}$ such that $k+1+i < \lambda, \langle \mathbf{H}, \mathbf{T} \rangle, k+1+i \models \psi$ and for all $0 \le j < i, \langle \mathbf{H}, \mathbf{T} \rangle, k+1+j \models \varphi$. From this and $\langle \mathbf{H}, \mathbf{T} \rangle, k+i' \models \psi$ and for all $0 \le j' < i, \langle \mathbf{H}, \mathbf{T} \rangle, k+j' \models \varphi$. Therefore, $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi \bigcup_n \psi$.

• For $\varphi \mathbb{R}_n \psi \leftrightarrow \psi \land (\varphi \lor \widehat{\circ} \varphi \mathbb{R}_{n-1} \psi)$, let us consider first the left to right direction. For this, let us assume for the sake of contradiction that $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \psi \land (\varphi \lor \widehat{\circ} (\varphi \mathbb{R}_{n-1} \psi))$. If $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \psi$ then $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \mathbb{R}_n \psi$ by definition (just take i = 0). If $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \land \widehat{\circ} (\varphi \mathbb{R}_{n-1} \psi)$ then $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi, k+1 < \lambda$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k+1 \not\models \varphi \mathbb{R}_{n-1} \psi$. This means that there exists $0 \leq i < \overline{n-1}$ such that $k+1+i < \lambda, \langle \mathbf{H}, \mathbf{T} \rangle, k+1+i \not\models \psi$ and for all $0 \leq j < i, \langle \mathbf{H}, \mathbf{T} \rangle, k+1+j \not\models \varphi$. From this and $\langle \mathbf{H}, \mathbf{T} \rangle, k+i' \not\models \psi$ and for all $0 \leq j' < i, \langle \mathbf{H}, \mathbf{T} \rangle, k+j' \not\models \varphi$. Therefore, $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \mathbb{R}_n \psi$: a contradiction.

From right to left, let us assume by contradiction that $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \mathbb{R}_n \psi$. This means that there exists $0 \leq i < \overline{n}$ such that $k + i < \lambda$, $\langle \mathbf{H}, \mathbf{T} \rangle, k + i \not\models \psi$ and for all $0 \leq j < i, \langle \mathbf{H}, \mathbf{T} \rangle, k + j \not\models \varphi$. If i = 0 then we can easily conclude $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \psi$. If i > 0 then $\langle \mathbf{H}, \mathbf{T} \rangle, k + 1 \not\models \varphi \mathbb{R}_{n-1} \psi$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi$. As a consequence, $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \psi \land (\varphi \lor \widehat{0} (\varphi \mathbb{R}_{n-1} \psi))$.

• For $\varphi \mathbf{S}_n \psi \leftrightarrow \psi \lor (\varphi \land \bullet \varphi \mathbf{S}_{n-1} \psi)$, let us assume that $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi \mathbf{S}_n \psi$. This means that there exists $0 \leq i < \overline{n}$ such that $k - i \geq 0$, $\langle \mathbf{H}, \mathbf{T} \rangle, k - i \models \psi$ and for all $0 \leq j < i, \langle \mathbf{H}, \mathbf{T} \rangle, k - j \models \varphi$. If i = 0 then we can easily conclude $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \psi$. If i > 0 then $\langle \mathbf{H}, \mathbf{T} \rangle, k - 1 \models \varphi \mathbf{S}_{n-1} \psi$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi$. As a consequence, $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \psi \lor (\varphi \land \bullet (\varphi \mathbf{S}_{n-1} \psi))$.

For the converse direction, assume that $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \psi \lor (\varphi \land \bullet (\varphi \mathbb{U}_{n-1}\psi))$. If $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \psi$ then $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi \mathbf{S}_n \psi$ by definition (just take i = 0). If $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi \land \bullet (\varphi \mathbf{S}_{n-1}\psi)$ then $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi, k-1 \ge 0$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k-1 \models \varphi \mathbf{S}_{n-1}\psi$. This means that there exists $0 \le i < \overline{n-1}$ such that $k-1-i \ge 0, \langle \mathbf{H}, \mathbf{T} \rangle, k-1-i \models \psi$ and for all $0 \le j < i, \langle \mathbf{H}, \mathbf{T} \rangle, k-1-j \models \varphi$. From this and $\langle \mathbf{H}, \mathbf{T} \rangle, k-i' \models \psi$ and for all $0 \le j' < i, \langle \mathbf{H}, \mathbf{T} \rangle, k-j' \models \varphi$. Therefore, $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi \mathbf{S}_n \psi$.

• For $\varphi \mathbf{T}_n \psi \leftrightarrow \psi \land (\varphi \lor \widehat{\mathbf{0}} \varphi \mathbf{T}_{n-1} \psi)$, let us consider first the left to right direction. For this, let us assume for the sake of contradiction that $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \psi \land (\varphi \lor \widehat{\mathbf{0}} (\varphi \mathbf{T}_{n-1} \psi))$. If $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \psi$ then $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \mathbf{T}_n \psi$ by definition (just take i = 0). If $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \land \widehat{\mathbf{0}} (\varphi \mathbf{T}_{n-1} \psi)$ then $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi, k > 0$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k - 1 \not\models \varphi \mathbf{T}_{n-1} \psi$. This means that there exists $0 \leq i < \overline{n-1}$ such that

 $k-1-i \ge 0, \langle \mathbf{H}, \mathbf{T} \rangle, k-1-i \not\models \psi$ and for all $0 \le j < i, \langle \mathbf{H}, \mathbf{T} \rangle, k-1-j \not\models \varphi$. From this and $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi$ we conclude that there exists $0 \le i' < \overline{n}$ such that $k-i' \ge 0$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k-i' \not\models \psi$ and for all $0 \le j' < i, \langle \mathbf{H}, \mathbf{T} \rangle, k-j' \not\models \varphi$. Therefore, $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \mathbf{T}_n \psi$: a contradiction.

From right to left, let us assume by contradiction that $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \mathbf{T}_n \psi$. This means that there exists $0 \leq i < \overline{n}$ such that $k - i \geq 0$, $\langle \mathbf{H}, \mathbf{T} \rangle, k - i \not\models \psi$ and for all $0 \leq j < i, \langle \mathbf{H}, \mathbf{T} \rangle, k - j \not\models \varphi$. If i = 0 then we can easily conclude $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \psi$. If i > 0 then $\langle \mathbf{H}, \mathbf{T} \rangle, k - 1 \not\models \varphi \mathbf{T}_{n-1} \psi$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi$. As a consequence, $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \psi \land (\varphi \lor \widehat{\bullet} (\varphi \mathbf{T}_{n-1} \psi))$.

QED

Proof of Proposition 7. We consider the different cases below:

• For $\varphi \mathbb{U}_{\ell} \psi \leftrightarrow \psi \lor (\varphi \land \circ \varphi \mathbb{U}_{\ell} \psi)$, let us consider from left to right that $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi \mathbb{U}_{\ell} \psi$. This means that there exists $0 \leq i < \lambda$ such that $k + i < \lambda, \langle \mathbf{H}, \mathbf{T} \rangle, k + i \models \psi$ and for all $0 \leq j < i, \langle \mathbf{H}, \mathbf{T} \rangle, k + j \models \varphi$. If i = 0 then we can easily conclude $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \psi$. If $0 < i < \lambda$ then $\langle \mathbf{H}, \mathbf{T} \rangle, k + 1 \models \varphi \mathbb{U}_{\ell} \psi$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi$. As a consequence, $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \psi \lor (\varphi \land \circ (\varphi \mathbb{U}_{\ell} \psi))$.

For the converse direction, assume that $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \psi \lor (\varphi \land \circ (\varphi \boxtimes_{\ell} \psi))$. If $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \psi$ then $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi \boxtimes_{\ell} \psi$ by definition (just take i = 0). If $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi \land \circ (\varphi \boxtimes_{\ell} \psi)$ then $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi, k + 1 < \lambda$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k + 1 \models \varphi \boxtimes_{\ell} \psi$. This means that there exists $0 \le i < \lambda$ such that $k + 1 + i < \lambda, \langle \mathbf{H}, \mathbf{T} \rangle, k + 1 + i \models \psi$ and for all $0 \le j < i$, $\langle \mathbf{H}, \mathbf{T} \rangle, k + 1 + j \models \varphi$. From this and $\langle \mathbf{H}, \mathbf{T} \rangle, k + i' \models \psi$ and for all $0 \le j' < i$, $\langle \mathbf{H}, \mathbf{T} \rangle, k + j' \models \varphi$. Therefore, $\langle \mathbf{H}, \mathbf{T} \rangle, k + i' \models \psi$ and for all $0 \le j' < i$, $\langle \mathbf{H}, \mathbf{T} \rangle, k + j' \models \varphi$. Therefore, $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi \boxtimes_{\ell} \psi$.

• For $\varphi \mathbb{R}_{\ell} \psi \leftrightarrow \psi \land (\varphi \lor \widehat{\circ} \varphi \mathbb{R}_{\ell} \psi)$, let us consider first the left to right direction. For this, let us assume for the sake of contradiction that $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \psi \land (\varphi \lor \widehat{\circ} (\varphi \mathbb{R}_{\ell} \psi))$. If $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \psi$ then $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \mathbb{R}_{\ell} \psi$ by definition (just take i = 0). If $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \lor \widehat{\circ} (\varphi \mathbb{R}_{\ell} \psi)$ then $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi, k + 1 < \lambda$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k + 1 \not\models \varphi \mathbb{R}_{\ell} \psi$. This means that there exists $0 \leq i < \lambda$ such that $k + 1 + i < \lambda$, $\langle \mathbf{H}, \mathbf{T} \rangle, k + 1 + i \not\models \psi$ and for all $0 \leq j < i$, $\langle \mathbf{H}, \mathbf{T} \rangle, k + 1 + j \not\models \varphi$. From this and $\langle \mathbf{H}, \mathbf{T} \rangle, k \neq \varphi$ we conclude that there exists $0 \leq i' < \lambda$ such that $k + i' < \lambda$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k + i' \not\models \psi$ and for all $0 \leq j' < i$, $\langle \mathbf{H}, \mathbf{T} \rangle, k + j' \not\models \varphi$. Therefore, $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \mathbb{R}_{\ell} \psi$: a contradiction.

From right to left, let us assume by contradiction that $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \mathbb{R}_{\ell} \psi$. This means that there exists $0 \leq i < \lambda$ such that $k + i < \lambda$, $\langle \mathbf{H}, \mathbf{T} \rangle, k + i \not\models \psi$ and for all $0 \leq j < i$, $\langle \mathbf{H}, \mathbf{T} \rangle, k + j \not\models \varphi$. If i = 0 then we can easily conclude $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \psi$. If i > 0 then $\langle \mathbf{H}, \mathbf{T} \rangle, k + 1 \not\models \varphi \mathbb{R}_{\ell} \psi$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi$. As a consequence, $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \psi \land (\varphi \lor \widehat{\circ} (\varphi \mathbb{R}_{\ell} \psi))$.

• For $\varphi \mathbf{S}_{\boldsymbol{\ell}} \psi \leftrightarrow \psi \lor (\varphi \land \mathbf{\bullet} \varphi \mathbf{S}_{\boldsymbol{\ell}} \psi)$, let us assume that $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi \mathbf{S}_{\boldsymbol{\ell}} \psi$. This means that there exists $0 \leq i < \lambda$ such that $k - i \geq 0$, $\langle \mathbf{H}, \mathbf{T} \rangle, k - i \models \psi$ and for all $0 \leq j < i, \langle \mathbf{H}, \mathbf{T} \rangle, k - j \models \varphi$. If i = 0 then we can easily conclude $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \psi$. If i > 0 then $\langle \mathbf{H}, \mathbf{T} \rangle, k - 1 \models \varphi \mathbf{S}_{\boldsymbol{\ell}} \psi$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi$. As a consequence, $\langle \mathbf{H}, \mathbf{T} \rangle, k \models$ $\psi \lor (\varphi \land \mathbf{\bullet} (\varphi \mathbf{S}_{\boldsymbol{\ell}} \psi))$.

For the converse direction, assume that $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \psi \lor (\varphi \land \bullet (\varphi \cup_{\ell} \psi))$. If $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \psi$ then $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi \land \bullet (\varphi \mathbf{S}_{\ell} \psi)$ by definition (just take i = 0). If $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi \land \bullet (\varphi \mathbf{S}_{\lambda} \psi)$ then $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi, k - 1 \ge 0$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k - 1 \models \varphi \mathbf{S}_n \psi$. This means that there exists $0 \le i < \lambda$ such that $k - 1 - i \ge 0$, $\langle \mathbf{H}, \mathbf{T} \rangle, k - 1 = \psi$ and for all $0 \le j < i$,

 $\langle \mathbf{H}, \mathbf{T} \rangle, k-1-j \models \varphi$. From this and $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi$ we conclude that there exists $0 \leq i' < \lambda$ such that $k-i' \geq 0$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k-i' \models \psi$ and for all $0 \leq j' < i$, $\langle \mathbf{H}, \mathbf{T} \rangle, k-j' \models \varphi$. Therefore, $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi \mathbf{S}_{\ell} \psi$.

• For $\varphi \mathbf{T}_{\boldsymbol{\ell}} \psi \leftrightarrow \psi \land (\varphi \lor \widehat{\mathbf{o}} \varphi \mathbf{T}_{\lambda} \psi)$, let us consider first the left to right direction. For this, let us assume for the sake of contradiction that $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \psi \land (\varphi \lor \widehat{\mathbf{o}} (\varphi \mathbf{T}_{\lambda} \psi))$. If $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \psi$ then $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \mathbf{T}_{\lambda} \psi$ by definition (just take i = 0). If $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \land \widehat{\mathbf{o}} (\varphi \mathbf{T}_{\lambda} \psi)$ then $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi, k > 0$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k - 1 \not\models \varphi \mathbf{T}_{\boldsymbol{\ell}} \psi$. This means that there exists $0 \leq i < \lambda$ such that $k - 1 - i \geq 0$, $\langle \mathbf{H}, \mathbf{T} \rangle, k - 1 - i \not\models \psi$ and for all $0 \leq j < i$, $\langle \mathbf{H}, \mathbf{T} \rangle, k - 1 - j \not\models \varphi$. From this and $\langle \mathbf{H}, \mathbf{T} \rangle, k - i' \not\models \psi$ and for all $0 \leq j' < i$, $\langle \mathbf{H}, \mathbf{T} \rangle, k - j' \not\models \varphi$. Therefore, $\langle \mathbf{H}, \mathbf{T} \rangle, k \neq \sigma \mathbf{T}_{\boldsymbol{\ell}} \psi$: a contradiction. From right to left, let us assume by contradiction that $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \mathbf{T}_{\boldsymbol{\ell}} \psi$. This means that there exists $0 \leq i < \lambda$ such that k - i > 0, $\langle \mathbf{H}, \mathbf{T} \rangle, k - i \not\models \psi$ and

means that there exists $0 \leq i < \lambda$ such that $k - i \geq 0$, $\langle \mathbf{H}, \mathbf{T} \rangle$, $k - i \not\models \psi$ and for all $0 \leq j < i$, $\langle \mathbf{H}, \mathbf{T} \rangle$, $k - j \not\models \varphi$. If i = 0 then we can easily conclude $\langle \mathbf{H}, \mathbf{T} \rangle$, $k \not\models \psi$. If i > 0 then $\langle \mathbf{H}, \mathbf{T} \rangle$, $k - 1 \not\models \varphi \mathbf{T}_{\ell} \psi$ and $\langle \mathbf{H}, \mathbf{T} \rangle$, $k \not\models \varphi$. As a consequence, $\langle \mathbf{H}, \mathbf{T} \rangle$, $k \not\models \psi \land (\varphi \lor \widehat{\bullet} (\varphi \mathbf{T}_{\ell} \psi))$.

QED

Proof of Lemma 12. The proof is done by using double induction on the complexity of the formula and on \overline{n} . We consider the metric operators below

- Case $\varphi \mathbb{U}_1 \psi$: from left to right, if $\mathbf{M}, k \models \varphi \mathbb{U}_1 \psi$ in MHT then, by Proposition 6, $\mathbf{M}, k \models \psi$ in MHT. By induction hypothesis, $\mathbf{M}, k \models \tau(\psi)$ in THT and, consequently, $\mathbf{M}, k \models \tau(\varphi \mathbb{U}_1 \psi)$ in THT. Conversely, if $\mathbf{M}, k \models \tau(\varphi \mathbb{U}_1 \psi)$ in THT then, by definition, $\mathbf{M}, k \models \tau(\psi)$ in THT and, by induction hypothesis, $\mathbf{M}, k \models \psi$ in MHT and, thanks to Proposition 6 we conclude $\mathbf{M}, k \models \varphi \mathbb{U}_1 \psi$.
- Cases $\varphi \mathbb{R}_1 \psi$, $\varphi \mathbf{S}_1 \psi$ and $\varphi \mathbf{T}_1 \psi$ are proved in a similar way.
- Case $\varphi \mathbb{U}_n \psi$ with n > 1: from left to right, from $\mathbf{M}, k \models \varphi \mathbb{U}_n \psi$ and Proposition 6, $\mathbf{M}, k \models \psi \lor (\varphi \land \circ (\varphi \mathbb{U}_{n-1} \psi))$ in MHT. By using induction on the subformulas and on \overline{n} we can conclude that $\mathbf{M}, k \models \tau(\psi)$ in THT or both $\mathbf{M}, k \models \tau(\varphi)$ and $\mathbf{M}, k \models \circ \tau(\varphi \mathbb{U}_{n-1} \psi)$ in THT. By using the definition of $\tau()$ it follows that $\mathbf{M}, k \models$ $\tau(\psi \lor (\varphi \land \circ (\varphi \mathbb{U}_{n-1} \psi)))$ and so $\mathbf{M}, k \models \tau(\varphi \mathbb{U}_n \psi)$ in THT. Conversely, if $\mathbf{M}, k \models$ $\tau(\varphi \mathbb{U}_n \psi)$ in THT then, by using the definition of $\tau()$, we conclude that $\mathbf{M}, k \models$ $\tau(\psi) \lor (\tau(\varphi) \land \circ \tau(\varphi \mathbb{U}_{n-1} \psi))$ in THT. By applying induction on \overline{n} and on the subformulas we can easily conclude that $\mathbf{M}, k \models \psi \lor (\varphi \land \circ (\varphi \mathbb{U}_{n-1} \psi))$ in MHT. Thanks to Proposition 6 we conclude that $\mathbf{M}, k \models \varphi \mathbb{U}_n \psi$ in MHT.
- Case $\varphi \mathbb{R}_n \psi$ with $\overline{n} > 1$: from left to right, assume by contradiction that $\mathbf{M}, k \not\models \tau(\varphi \mathbb{R}_n \psi)$ in THT. By definition of $\tau(\cdot)$ we can derive that $\mathbf{M}, k \not\models \tau(\psi) \land (\tau(\varphi) \lor \widehat{\circ} \tau(\varphi \mathbb{R}_{n-1}\psi))$ in THT. By applying induction on \overline{n} and on the subformulas we can easily conclude that $\mathbf{M}, k \not\models \psi \land (\varphi \lor \widehat{\circ} (\varphi \mathbb{R}_{n-1}\psi))$ in MHT. Thanks to Proposition 6 we conclude that $\mathbf{M}, k \not\models \varphi \mathbb{R}_n \psi$ in MHT: a contradiction. For the converse direction assume, again, by contradiction that $\mathbf{M}, k \not\models \varphi \mathbb{R}_n \psi$ and Proposition 6, $\mathbf{M}, k \not\models \psi \land (\varphi \lor \widehat{\circ} (\varphi \mathbb{R}_{n-1}\psi))$ in MHT. By using induction on the subformulas and on \overline{n} we can conclude that $\mathbf{M}, k \not\models \tau(\psi)$ in THT or both $\mathbf{M}, k \not\models \tau(\varphi)$ and $\mathbf{M}, k \not\models \widehat{\circ} \tau(\varphi \mathbb{R}_{n-1}\psi)$ in THT. By using the definition of $\tau(\cdot)$ it follows that $\mathbf{M}, k \not\models \tau(\psi \land (\varphi \lor \widehat{\circ} (\varphi \mathbb{R}_{n-1}\psi)))$ and so $\mathbf{M}, k \not\models \tau(\varphi \mathbb{R}_n\psi)$ in THT: a contradiction.
- Case $\varphi \mathbf{S}_n \psi$ with $\overline{n} > 1$: from left to right, from $\mathbf{M}, k \models \varphi \mathbf{S}_n \psi$ and Proposition 6,

20

 $\mathbf{M}, k \models \psi \lor (\varphi \land \bullet (\varphi \mathbf{S}_{n-1}\psi))$ in MHT. By using induction on the subformulas and on \overline{n} we can conclude that $\mathbf{M}, k \models \tau(\psi)$ in THT or both $\mathbf{M}, k \models \tau(\varphi)$ and $\mathbf{M}, k \models \bullet \tau(\varphi \mathbf{S}_{n-1}\psi)$ in THT. By using the definition of $\tau()$ it follows that $\mathbf{M}, k \models$ $\tau(\psi \lor (\varphi \land \bullet (\varphi \mathbf{S}_{n-1}\psi)))$ and so $\mathbf{M}, k \models \tau(\varphi \mathbf{S}_n\psi)$ in THT. Conversely, if $\mathbf{M}, k \models$ $\tau(\varphi \mathbf{S}_n\psi)$ in THT then, by using the definition of $\tau()$, we conclude that $\mathbf{M}, k \models$ $\tau(\psi) \lor (\tau(\varphi) \land \bullet \tau(\varphi \mathbf{S}_{n-1}\psi))$ in THT. By applying induction on \overline{n} and on the subformulas we can easily conclude that $\mathbf{M}, k \models \psi \lor (\varphi \land \bullet (\varphi \mathbf{S}_{n-1}\psi))$ in MHT. Thanks to Proposition 6 we conclude that $\mathbf{M}, k \models \varphi \mathbf{S}_n\psi$ in MHT.

- Case $\varphi \mathbf{T}_n \psi$ with $\overline{n} > 1$: from left to right, assume by contradiction that $\mathbf{M}, k \not\models \tau(\varphi \mathbf{T}_n \psi)$ in THT. By definition of $\tau(\cdot)$ we can derive that $\mathbf{M}, k \not\models \tau(\psi) \wedge (\tau(\varphi) \vee \hat{\mathbf{o}} \tau(\varphi \mathbf{T}_{n-1}\psi))$ in THT. By applying induction on \overline{n} and on the subformulas we can easily conclude that $\mathbf{M}, k \not\models \psi \wedge (\varphi \vee \hat{\mathbf{o}} (\varphi \mathbf{T}_{n-1}\psi))$ in MHT. Thanks to Proposition 6 we conclude that $\mathbf{M}, k \not\models \varphi \mathbf{T}_n \psi$ in MHT: a contradiction. For the converse direction assume, again, by contradiction that $\mathbf{M}, k \not\models \varphi \mathbf{T}_n \psi$ and Proposition 6, $\mathbf{M}, k \not\models \psi \wedge (\varphi \vee \hat{\mathbf{o}} (\varphi \mathbf{T}_{n-1}\psi))$ in MHT. By using induction on the subformulas and on \overline{n} we can conclude that $\mathbf{M}, k \not\models \tau(\psi)$ in THT or both $\mathbf{M}, k \not\models \tau(\varphi)$ and $\mathbf{M}, k \not\models \hat{\mathbf{o}} \tau(\varphi \mathbf{T}_{n-1}\psi)$ in THT. By using the definition of $\tau(\cdot)$ it follows that $\mathbf{M}, k \not\models \tau(\psi \wedge (\varphi \vee \hat{\mathbf{o}} (\varphi \mathbf{T}_{n-1}\psi)))$ and so $\mathbf{M}, k \not\models \tau(\varphi \mathbf{T}_n\psi)$ in THT: a contradiction.
- Case $\varphi \mathbb{U}_{\ell} \psi$: from left to right, from $\mathbf{M}, k \models \varphi \mathbb{U}_{\ell} \psi$ and Proposition 7, $\mathbf{M}, k \models \psi \lor (\varphi \land \circ (\varphi \mathbb{U}_{\ell} \psi))$ in MHT. By using induction on the subformulas we can conclude that $\mathbf{M}, k \models \tau(\psi)$ in THT or both $\mathbf{M}, k \models \tau(\varphi)$ and $\mathbf{M}, k \models \circ \tau(\varphi \mathbb{U}_{\ell} \psi)$ in THT. By using the definition of $\tau()$ it follows that $\mathbf{M}, k \models \tau(\psi \lor (\varphi \land \circ (\varphi \mathbb{U}_{\ell} \psi)))$ and so $\mathbf{M}, k \models \tau(\varphi \mathbb{U}_{\ell} \psi)$ in THT. Conversely, if $\mathbf{M}, k \models \tau(\varphi \mathbb{U}_{\ell} \psi)$ in THT then, by using the definition of $\tau()$, we conclude that $\mathbf{M}, k \models \tau(\psi) \lor (\tau(\varphi) \land \circ \tau(\varphi \mathbb{U}_{\ell} \psi))$ in THT. By applying induction on the subformulas we can easily conclude that $\mathbf{M}, k \models \psi \lor (\varphi \land \circ (\varphi \mathbb{U}_{\ell} \psi))$ in MHT. Thanks to Proposition 7 we conclude that $\mathbf{M}, k \models \varphi \mathbb{U}_{\ell} \psi$ in MHT.
- Case $\varphi \mathbb{R}_{\ell} \psi$: from left to right, assume by contradiction that $\mathbf{M}, k \not\models \tau(\varphi \mathbb{R}_{\ell} \psi)$ in THT. By definition of $\tau(\cdot)$ we can derive that $\mathbf{M}, k \not\models \tau(\psi) \land (\tau(\varphi) \lor \widehat{\circ} \tau(\varphi \mathbb{R}_{\ell} \psi))$ in THT. By applying induction on the subformulas we can easily conclude that $\mathbf{M}, k \not\models \psi \land (\varphi \lor \widehat{\circ} (\varphi \mathbb{R}_{\ell} \psi))$ in MHT. Thanks to Proposition 7 we conclude that $\mathbf{M}, k \not\models \varphi \mathbb{R}_{\ell} \psi$ in MHT: a contradiction. For the converse direction assume, again, by contradiction that $\mathbf{M}, k \not\models \varphi \mathbb{R}_{\ell} \psi$ and Proposition 7, $\mathbf{M}, k \not\models \psi \land (\varphi \lor \widehat{\circ} (\varphi \mathbb{R}_{\ell} \psi))$ in MHT. By using induction on the subformulas we can conclude that $\mathbf{M}, k \not\models \tau(\psi)$ in THT or both $\mathbf{M}, k \not\models \tau(\varphi)$ and $\mathbf{M}, k \not\models \widehat{\circ} \tau(\varphi \mathbb{R}_{\ell} \psi)$ in THT. By using the definition of $\tau(\cdot)$ it follows that $\mathbf{M}, k \not\models \tau(\psi \land (\varphi \lor \widehat{\circ} (\varphi \mathbb{R}_{\ell} \psi)))$ and so $\mathbf{M}, k \not\models \tau(\varphi \mathbb{R}_{\ell} \psi)$ in THT: a contradiction.
- Case $\varphi \mathbf{S}_{\boldsymbol{\ell}} \psi$: from left to right, from $\mathbf{M}, k \models \varphi \mathbf{S}_{\boldsymbol{\ell}} \psi$ and Proposition 7, $\mathbf{M}, k \models \psi \lor (\varphi \land \bullet (\varphi \mathbf{S}_{\boldsymbol{\ell}} \psi))$ in MHT. By using induction on the subformulas we can conclude that $\mathbf{M}, k \models \tau(\psi)$ in THT or both $\mathbf{M}, k \models \tau(\varphi)$ and $\mathbf{M}, k \models \bullet \tau(\varphi \mathbf{S}_{\boldsymbol{\ell}} \psi)$ in THT. By using the definition of $\tau()$ it follows that $\mathbf{M}, k \models \tau(\psi \lor (\varphi \land \bullet (\varphi \mathbf{S}_{\boldsymbol{\ell}} \psi)))$ and so $\mathbf{M}, k \models \tau(\varphi \mathbf{S}_{\boldsymbol{\ell}} \psi)$ in THT. Conversely, if $\mathbf{M}, k \models \tau(\varphi \mathsf{S}_{\boldsymbol{\ell}} \psi)$ in THT then, by using the definition of $\tau()$, we conclude that $\mathbf{M}, k \models \tau(\varphi) \lor (\tau(\varphi) \land \bullet \tau(\varphi \mathbf{S}_{\boldsymbol{\ell}} \psi))$ in THT. By applying induction on the subformulas we can easily conclude that $\mathbf{M}, k \models \psi \lor (\varphi \land \bullet (\varphi \mathbf{S}_{\boldsymbol{\ell}} \psi))$ in MHT. Thanks to Proposition 7 we conclude that $\mathbf{M}, k \models \varphi \mathbf{S}_{\boldsymbol{\ell}} \psi$ in MHT.

• Case $\varphi \mathbf{T}_{\boldsymbol{\ell}} \psi$: from left to right, assume by contradiction that $\mathbf{M}, k \not\models \tau(\varphi \mathbf{T}_{\boldsymbol{\ell}} \psi)$ in THT. By definition of $\tau()$ we can derive that $\mathbf{M}, k \not\models \tau(\psi) \land (\tau(\varphi) \lor \widehat{\bullet} \tau(\varphi \mathbf{T}_{\boldsymbol{\ell}} \psi))$ in THT. By applying induction on the subformulas we can easily conclude that $\mathbf{M}, k \not\models \psi \land (\varphi \lor \widehat{\bullet} (\varphi \mathbf{T}_{\boldsymbol{\ell}} \psi))$ in MHT. Thanks to Proposition 7 we conclude that $\mathbf{M}, k \not\models \varphi \mathbf{T}_{\boldsymbol{\ell}} \psi$ in MHT: a contradiction. For the converse direction assume, again, by contradiction that $\mathbf{M}, k \not\models \varphi \mathbf{T}_{\boldsymbol{\ell}} \psi$ and Proposition 7, $\mathbf{M}, k \not\models \psi \land (\varphi \lor \widehat{\bullet} (\varphi \mathbf{T}_{\boldsymbol{\ell}} \psi))$ in MHT. By using induction on the subformulas we can conclude that $\mathbf{M}, k \not\models \tau(\psi)$ in THT or both $\mathbf{M}, k \not\models \tau(\varphi)$ and $\mathbf{M}, k \not\models \widehat{\bullet} \tau(\varphi \mathbf{T}_{\boldsymbol{\ell}} \psi)$ in THT. By using the definition of $\tau()$ it follows that $\mathbf{M}, k \not\models \tau(\psi \land (\varphi \lor \widehat{\bullet} (\varphi \mathbf{T}_{\boldsymbol{\ell}} \psi)))$ and so $\mathbf{M}, k \not\models \tau(\varphi \mathbf{T}_{\boldsymbol{\ell}} \psi)$ in THT: a contradiction.

The proof is done by structural induction. For each case, the proof can be done by using the equivalences of Proposition 6 and Proposition 7 together with some propositional reasoning. QED

Proof of Proposition 9. The proof is by structural induction. We consider only until/release and since/trigger.

- $\varphi \mathbb{U}_n \psi$: from left to right assume that $\langle \mathbf{H}, \mathbf{T} \rangle$, $k \models \varphi \mathbb{U}_n \psi$. Therefore, there exists $i \in [0..\overline{n})$ such that $k+i < \lambda$, $\langle \mathbf{H}, \mathbf{T} \rangle$, $k+i \models \psi$ and for all $j \in [0..i)$, $\langle \mathbf{H}, \mathbf{T} \rangle$, $k+j \models \varphi$. By induction hypothesis, $\mathbf{m}(k+i,\psi) = 2$ and $\mathbf{m}(k+j,\varphi) = 2$, for all $j \in [0..i)$. Therefore, $\min\{\mathbf{m}(k+i,\psi), \mathbf{m}(k+j,\varphi) \mid j \in [0..i), k+i < \lambda\} = 2$ and so $\max\{\min\{\mathbf{m}(k+i,\psi), \mathbf{m}(k+j,\varphi) \mid j \in [0..i), k+i < \lambda\} \mid i \in [0..\overline{n})\} = 2$. Conversely, assume that $\mathbf{m}(k,\varphi \mathbb{U}_n \psi) = \max\{\min\{\mathbf{m}(k+i,\psi), \mathbf{m}(k+j,\varphi) \mid j \in [0..i), k+i < \lambda\} \mid i \in [0..\overline{n})\} = 2$. Therefore there exists $0 \le i < \overline{n}$ such that $\min\{\mathbf{m}(k+i,\psi), \mathbf{m}(k+j,\varphi) \mid j \in [0..i), k+i < \lambda\} = 2$. As a consequence, $k+i < \lambda$, $\mathbf{m}(k+i,\psi) = 2$ and $\mathbf{m}(k+j,\varphi) = 2$, for all $0 \le j < i$. By induction induction it follows that $\langle \mathbf{H}, \mathbf{T} \rangle$, $k+i \models \psi$ and $\langle \mathbf{H}, \mathbf{T} \rangle$, $k \models \varphi \mathbb{U}_n \psi$. The second item is proved in a similar way.
- $\varphi \mathbb{R}_n \psi$: from left to right, assume that $\mathbf{m}(k, \varphi \mathbb{R}_n \psi) = \min\{\max\{\mathbf{m}(k+i, \psi), \mathbf{m}(k+j, \varphi) \mid j \in [0..i), k+i < \lambda\} \mid i \in [0..\overline{n})\} \neq 2$. therefore, there exists $0 \le i < \overline{n}$ such that $\max\{\mathbf{m}(k+i, \psi), \mathbf{m}(k+j, \varphi) \mid j \in [0..i), k+i < \lambda\} \neq 2$. This means that $k+i < \lambda$, $\mathbf{m}(k+i, \psi) \ne 2$ and $\mathbf{m}(k+j, \varphi) \ne 2$ for all $0 \le j < i$. By induction hypothesis, $\langle \mathbf{H}, \mathbf{T} \rangle, k+i \not\models \psi$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k+j \not\models \varphi$, for all $0 \le j < i$. From this we conclude that $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \mathbb{R}_n \psi$. Conversely, assume that $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \mathbb{R}_n \psi$. This means that there exists $i \in [0..\overline{n}]$ such that $k+i < \lambda$, $\langle \mathbf{H}, \mathbf{T} \rangle, k+i \not\models \psi$ and for all $j \in [0..i)$, $\langle \mathbf{H}, \mathbf{T} \rangle, k+j \not\models \varphi$. By induction hypothesis, $\mathbf{m}(k+i,\psi) \ne 2$ and $\mathbf{m}(k+j,\varphi) \ne 2$, for all $j \in [0..i)$. Therefore, $\max\{\mathbf{m}(k+i,\psi), \mathbf{m}(k+j,\varphi) \mid j \in [0..i), k+i < \lambda\} \ne 2$ and so $\min\{\max\{\mathbf{m}(k+i,\psi), \mathbf{m}(k+j,\varphi) \mid j \in [0..i), k+i < \lambda\} \mid i \in [0..\overline{n})\} = 2$: a contradiction. The second item is proved in a similar way.
- $\varphi \mathbf{S}_n \psi$ From left to right, assume that $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi \mathbf{S}_n \psi$. Therefore, there exists $i \in [0..\overline{n})$ such that $k-i \ge 0$, $\langle \mathbf{H}, \mathbf{T} \rangle, k-i \models \psi$ and for all $j \in [0..i)$, $\langle \mathbf{H}, \mathbf{T} \rangle, k-j \models \varphi$. By induction hypothesis, $\mathbf{m}(k-i,\psi) = 2$ and $\mathbf{m}(k-j,\varphi) = 2$, for all $j \in [0..i)$. Therefore, $\min\{\mathbf{m}(k-i,\psi), \mathbf{m}(k-j,\varphi) \mid j \in [0..i), k-i \ge 0\} = 2$ and so $\max\{\min\{\mathbf{m}(k-i,\psi), \mathbf{m}(k-j,\varphi) \mid j \in [0..i), k-i \ge 0\} \mid i \in [0..\overline{n})\} = 2$. Conversely, assume that $\mathbf{m}(k,\varphi \mathbf{S}_n \psi) = \max\{\min\{\mathbf{m}(k-i,\psi), \mathbf{m}(k-j,\varphi) \mid j \in [0..i), k-i \ge 0\} \mid i \in [0..\overline{n})\} = 2$. Therefore there exists $0 \le i < \overline{n}$ such that $\min\{\mathbf{m}(k-i,\psi), \mathbf{m}(k-j,\varphi) \mid j \in [0..i), k-i \ge 0\} = 2$. As a consequence, $k-i \ge 0$,

22

 $\boldsymbol{m}(k-i,\psi) = 2$ and $\boldsymbol{m}(k-j,\varphi) = 2$, for all $0 \le j < i$. By induction induction it follows that $\langle \mathbf{H}, \mathbf{T} \rangle, k-i \models \psi$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k-j \models \varphi$, for all $0 \le j < i$. This means that $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi \mathbf{S}_n \psi$. The second item is proved in a similar way.

- $\varphi \mathbf{T}_n \psi$: from left to right, assume that $\mathbf{m}(k, \varphi \mathbf{T}_n \psi) = \min\{\max\{\mathbf{m}(k-i,\psi), \mathbf{m}(k-j,\varphi) \mid j \in [0..i), k-i \geq 0\} \mid i \in [0..\overline{n})\} \neq 2$. therefore, there exists $0 \leq i < \overline{n}$ such that $\max\{\mathbf{m}(k-i,\psi), \mathbf{m}(k-j,\varphi) \mid j \in [0..i), k-i \geq 0\} \neq 2$. This means that $k-i \geq 0$, $\mathbf{m}(k-i,\psi) \neq 2$ and $\mathbf{m}(k-j,\varphi) \neq 2$ for all $0 \leq j < i$. By induction hypothesis, $\langle \mathbf{H}, \mathbf{T} \rangle, k-i \not\models \psi$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k-j \not\models \varphi$, for all $0 \leq j < i$. By induction hypothesis, $\langle \mathbf{H}, \mathbf{T} \rangle, k-i \not\models \psi$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k-j \not\models \varphi$, for all $0 \leq j < i$. By induction hypothesis, $\langle \mathbf{H}, \mathbf{T} \rangle, k-i \not\models \psi$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k-j \not\models \varphi$. Some that $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \mathbf{T}_n \psi$. This means that there exists $i \in [0..\overline{n})$ such that $k-i \geq 0$, $\langle \mathbf{H}, \mathbf{T} \rangle, k-i \not\models \psi$ and for all $j \in [0..i), \langle \mathbf{H}, \mathbf{T} \rangle, k-j \not\models \varphi$. By induction hypothesis, $\mathbf{m}(k-i,\psi) \neq 2$ and $\mathbf{m}(k-j,\varphi) \neq 2$, for all $j \in [0..i)$. Therefore, $\max\{\mathbf{m}(k-i,\psi), \mathbf{m}(k-j,\varphi) \mid j \in [0..i), k-i \geq 0\} \neq 2$ and so $\min\{\max\{\mathbf{m}(k-i,\psi), \mathbf{m}(k-j,\varphi) \mid j \in [0..i), k-i \geq 0\} \mid i \in [0..\overline{n})\} = 2$. The second item is proved in a similar way: a contradiction. The second item is proved in a similar way.
- $\varphi \bigcup_{\ell} \psi$: from left to right assume that $\langle \mathbf{H}, \mathbf{T} \rangle$, $k \models \varphi \bigcup_{\ell} \psi$. Therefore, there exists $i \in [0..\lambda)$ such that $k+i < \lambda$, $\langle \mathbf{H}, \mathbf{T} \rangle$, $k+i \models \psi$ and for all $j \in [0..i)$, $\langle \mathbf{H}, \mathbf{T} \rangle$, $k+j \models \varphi$. By induction hypothesis, $\mathbf{m}(k+i,\psi) = 2$ and $\mathbf{m}(k+j,\varphi) = 2$, for all $j \in [0..i)$. Therefore, $\min\{\mathbf{m}(k+i,\psi), \mathbf{m}(k+j,\varphi) \mid j \in [0..i), k+i < \lambda\} = 2$ and so $\max\{\min\{\mathbf{m}(k+i,\psi), \mathbf{m}(k+j,\varphi) \mid j \in [0..i), k+i < \lambda\} \mid i \in [0..\lambda)\} = 2$. Conversely, assume that $\mathbf{m}(k,\varphi \bigcup_{\ell} \psi) = \max\{\min\{\mathbf{m}(k+i,\psi), \mathbf{m}(k+j,\varphi) \mid j \in [0..i), k+i < \lambda\} \mid i \in [0..\lambda)\} = 2$. Therefore there exists $0 \le i < \lambda$ such that $\min\{\mathbf{m}(k+i,\psi), \mathbf{m}(k+j,\varphi) \mid j \in [0..i), k+i < \lambda\} = 2$. As a consequence, $k+i < \lambda$, $\mathbf{m}(k+i,\psi) = 2$ and $\mathbf{m}(k+j,\varphi) = 2$, for all $0 \le j < i$. By induction induction it follows that $\langle \mathbf{H}, \mathbf{T} \rangle$, $k \models \varphi \bigcup_{\ell} \psi$. The second item is proved in a similar way.
- $\varphi \mathbb{R}_{\ell} \psi$: from left to right, assume that $\mathbf{m}(k, \varphi \mathbb{R}_{\ell} \psi) = \min\{\max\{\mathbf{m}(k+i,\psi), \mathbf{m}(k+j,\varphi) \mid j \in [0..i), k+i < \lambda\} \mid i \in [0..\lambda)\} \neq 2$. therefore, there exists $0 \le i < \lambda$ such that $\max\{\mathbf{m}(k+i,\psi), \mathbf{m}(k+j,\varphi) \mid j \in [0..i), k+i < \lambda\} \neq 2$. This means that $k+i < \lambda, \mathbf{m}(k+i,\psi) \ne 2$ and $\mathbf{m}(k+j,\varphi) \ne 2$ for all $0 \le j < i$. By induction hypothesis, $\langle \mathbf{H}, \mathbf{T} \rangle, k+i \not\models \psi$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k+j \not\models \varphi$, for all $0 \le j < i$. From this we conclude that $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \mathbb{R}_{\ell} \psi$. Conversely, assume that $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \mathbb{R}_{\ell} \psi$. This means that there exists $i \in [0..\lambda)$ such that $k+i < \lambda, \langle \mathbf{H}, \mathbf{T} \rangle, k+i \not\models \psi$ and for all $j \in [0..i)$, $\langle \mathbf{H}, \mathbf{T} \rangle, k+j \not\models \varphi$. By induction hypothesis, $\mathbf{m}(k+i,\psi) \ne 2$ and $\mathbf{m}(k+j,\varphi) \ne 2$, for all $j \in [0..i)$. Therefore, $\max\{\mathbf{m}(k+i,\psi), \mathbf{m}(k+j,\varphi) \mid j \in [0..i), k+i < \lambda\} \neq 2$ and so $\min\{\max\{\mathbf{m}(k+i,\psi), \mathbf{m}(k+j,\varphi) \mid j \in [0..i), k+i < \lambda\} \mid i \in [0..\lambda)\} = 2$: a contradiction. The second item is proved in a similar way.
- $\varphi \mathbf{S}_{\boldsymbol{\ell}} \psi$ From left to right, assume that $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi \mathbf{S}_{\boldsymbol{\ell}} \psi$. Therefore, there exists $i \in [0..\lambda)$ such that $k-i \ge 0$, $\langle \mathbf{H}, \mathbf{T} \rangle, k-i \models \psi$ and for all $j \in [0..i)$, $\langle \mathbf{H}, \mathbf{T} \rangle, k-j \models \varphi$. By induction hypothesis, $\boldsymbol{m}(k-i,\psi) = 2$ and $\boldsymbol{m}(k-j,\varphi) = 2$, for all $j \in [0..i)$. Therefore, $\min\{\boldsymbol{m}(k-i,\psi), \boldsymbol{m}(k-j,\varphi) \mid j \in [0..i), k-i \ge 0\} = 2$ and so $\max\{\min\{\boldsymbol{m}(k-i,\psi), \boldsymbol{m}(k-j,\varphi) \mid j \in [0..i), k-i \ge 0\} \mid i \in [0..\lambda)\} = 2$. Conversely, assume that $\boldsymbol{m}(k,\varphi \mathbf{S}_{\boldsymbol{\ell}} \psi) = \max\{\min\{\boldsymbol{m}(k-i,\psi), \boldsymbol{m}(k-j,\varphi) \mid j \in [0..i), k-i \ge 0\} \mid i \in [0..\lambda)\} = 2$. Therefore there exists $0 \le i < \lambda$ such that $\min\{\boldsymbol{m}(k-i,\psi), \boldsymbol{m}(k-j,\varphi) \mid j \in [0..i), k-i \ge 0\} = 2$. As a consequence, $k-i \ge 0$,

 $m(k-i,\psi) = 2$ and $m(k-j,\varphi) = 2$, for all $0 \le j < i$. By induction it follows that $\langle \mathbf{H}, \mathbf{T} \rangle, k-i \models \psi$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k-j \models \varphi$, for all $0 \le j < i$. This means that $\langle \mathbf{H}, \mathbf{T} \rangle, k \models \varphi \mathbf{S}_{\ell} \psi$. The second item is proved in a similar way.

• $\varphi \mathbf{T}_{\boldsymbol{\ell}} \psi$: from left to right, assume that $\boldsymbol{m}(k, \varphi \mathbf{T}_{\boldsymbol{\ell}} \psi) = \min\{\max\{\boldsymbol{m}(k-i,\psi), \boldsymbol{m}(k-j,\varphi) \mid j \in [0..i), k-i \geq 0\} \mid i \in [0..\lambda)\} \neq 2$. therefore, there exists $0 \leq i < \lambda$ such that $\max\{\boldsymbol{m}(k-i,\psi), \boldsymbol{m}(k-j,\varphi) \mid j \in [0..i), k-i \geq 0\} \neq 2$. This means that $k-i \geq 0$, $\boldsymbol{m}(k-i,\psi) \neq 2$ and $\boldsymbol{m}(k-j,\varphi) \neq 2$ for all $0 \leq j < i$. By induction hypothesis, $\langle \mathbf{H}, \mathbf{T} \rangle, k-i \not\models \psi$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k-j \not\models \varphi$, for all $0 \leq j < i$. By induction hypothesis, $\langle \mathbf{H}, \mathbf{T} \rangle, k-i \not\models \psi$ and $\langle \mathbf{H}, \mathbf{T} \rangle, k-j \not\models \varphi$, for all $0 \leq j < i$. From this we conclude that $\langle \mathbf{H}, \mathbf{T} \rangle, k \not\models \varphi \mathbf{T}_{\boldsymbol{\ell}} \psi$: a contradiction. Conversely, assume that $\langle \mathbf{H}, \mathbf{T} \rangle, k-i \not\models \psi$ and for all $j \in [0..i), \langle \mathbf{H}, \mathbf{T} \rangle, k-j \not\models \varphi$. By induction hypothesis, $\boldsymbol{m}(k-i,\psi) \neq 2$ and $\boldsymbol{m}(k-j,\varphi) \neq 2$, for all $j \in [0..i)$. Therefore, $\max\{\boldsymbol{m}(k-i,\psi), \boldsymbol{m}(k-j,\varphi) \mid j \in [0..i), k-i \geq 0\} \neq 2$ and so $\min\{\max\{\boldsymbol{m}(k-i,\psi), \boldsymbol{m}(k-j,\varphi) \mid j \in [0..i), k-i \geq 0\} \mid i \in [0..\lambda)\} = 2$. The second item is proved in a similar way: a contradiction. The second item is proved in a similar way.

QED **Proof of Proposition 13.** We assume k_{φ} is defined for any metric formula φ as stated in the proposition. We proceed by structural induction.

- case φ = p: in this case |p| = 1 and cl(p) = {p}, so |cl(p)| = 1. Since there is no metric connective in p, k_p = 1. Therefore, |cl(p)| = 1 ≤ 2 * 1 * |p| = 2.
- case $\varphi \wedge \psi$: in this case, $\operatorname{cl}(\varphi \wedge \psi) = \{\varphi \wedge \psi\} \cup \operatorname{cl}(\varphi) \cup \operatorname{cl}(\psi)$ and $|\operatorname{cl}(\varphi \wedge \psi)| \leq 1 + |\operatorname{cl}(\varphi)| + |\operatorname{cl}(\psi)|$. We assume, without loss of generality, $\max(k_{\varphi}, k_{\psi}) = k_{\psi}$ when needed. We prove this case next:

$$\begin{aligned} \operatorname{cl}(\varphi \wedge \psi) &| \leq 1 + |\operatorname{cl}(\varphi)| + |\operatorname{cl}(\psi)| \\ &\leq 1 + 2k_{\varphi}|\varphi| + 2k_{\psi}|\psi| & \text{by induction on } \varphi \text{ and } \psi \\ &\leq 1 + 2k_{\varphi}|\varphi| + 2k_{\psi}|\psi| + 2(k_{\psi} - k_{\varphi})|\varphi| & 2(k_{\psi} - k_{\varphi})|\varphi| \geq 0 \\ &\leq 1 + 2k_{\varphi}|\varphi| + 2k_{\psi}|\psi| + 2k_{\psi}|\varphi| \\ &\leq 1 + 2k_{\psi}|\psi| + 2k_{\psi}|\varphi| \\ &\leq 1 + 2k_{\psi}(|\psi| + |\varphi|) \\ &\leq 1 + 2k_{\psi}(|\varphi \wedge \psi| - 1) \\ &\leq 1 - 2k_{\psi} + 2k_{\psi}|\varphi \wedge \psi| \\ &\leq 2k_{\psi}|\varphi \wedge \psi|. & 1 - 2k_{\psi} < 0. \end{aligned}$$

- the proof for the formulas $\varphi \lor \psi$ and $\varphi \to \psi$ is done as for $\varphi \land \psi$.
- case $\circ\varphi$: in this case, $\operatorname{cl}(\circ\varphi) = \{\circ\varphi\} \cup \operatorname{cl}(\varphi) \text{ and } |\circ\varphi| = 1 + |\varphi|$. It follows that

$$\begin{aligned} |\mathrm{cl}(\circ\varphi)| &\leq 1 + |\mathrm{cl}(\varphi)| \\ &\leq 1 + 2k_{\varphi}|\varphi| & \text{by induction on } \varphi \\ &\leq 1 + 2k_{\varphi} \left(|\circ\varphi| - 1\right) \\ &\leq 1 - 2k_{\varphi} + 2k_{\varphi}|\circ\varphi| \\ &\leq 2k_{\varphi}|\circ\varphi| & 1 - 2k_{\varphi} < 0 \end{aligned}$$

24

- The proof for $\widehat{\circ}\varphi$, $\bullet\varphi$ and $\widehat{\bullet}\varphi$ follows the same line of reasoning as for $\circ\varphi$.
- Case $\varphi \mathbb{U}_{\ell} \psi$: in this case, $\operatorname{cl}(\varphi \mathbb{U}_{\ell} \psi) = \{\varphi \mathbb{U}_{\ell} \psi, \circ (\varphi \mathbb{U}_{\ell} \psi)\} \cup \operatorname{cl}(\varphi) \cup \operatorname{cl}(\psi) \text{ and } |\varphi \mathbb{U}_{\ell} \psi| = 1 + |\varphi| + |\psi|$. We proceed in a similar way as in the previous cases. We will also assume, without loss of generality, that $\max(k_{\varphi}, k_{\psi}) = k_{\psi}$ when needed.

$$\begin{aligned} |\mathrm{cl}(\varphi \mathbb{U}_{\ell} \psi)| &\leq 2 + |\mathrm{cl}(\varphi)| + |\mathrm{cl}(\psi)| \\ &\leq 2 + 2k_{\varphi}|\varphi| + 2k_{\psi}|\psi| & \text{by induction on } \varphi \text{ and } \psi \\ &\leq 2 + 2k_{\varphi}|\varphi| + 2k_{\psi}|\psi| + 2(k_{\psi} - k_{\varphi})|\varphi| & 2(k_{\psi} - k_{\varphi})|\varphi| \geq 0 \\ &\leq 2 + 2k_{\varphi}|\varphi| + 2k_{\psi}|\psi| + 2k_{\psi}|\varphi| - 2k_{\varphi}|\varphi| \\ &\leq 2 + 2k_{\psi}|\psi| + 2k_{\psi}|\varphi| \\ &\leq 2 + 2k_{\psi}(|\psi| + |\varphi|) \\ &\leq 2 + 2k_{\psi}(|\varphi \mathbb{U}_{\ell} \psi| - 1) \\ &\leq 2 - 2k_{\psi} + 2k_{\psi}|\varphi \mathbb{U}_{\ell} \psi| \\ &\leq 2k_{\psi}|\varphi \mathbb{U}_{\ell} \psi|. & 2 - 2k_{\psi} \leq 0 \end{aligned}$$

- The proof for $\varphi \mathbb{R}_{\ell} \psi$, $\varphi \mathbf{S}_{\ell} \psi$ and $\varphi \mathbf{T}_{\ell} \psi$ follows the same reasoning as for the case $\varphi \mathbb{U}_{\ell} \psi$.
- Case $\varphi \mathbb{U}_n \psi$: in this case we have that $|\varphi \mathbb{U}_n \psi| = 1 + |\varphi| + |\psi|$ and $\operatorname{cl}(\varphi \mathbb{U}_n \psi) = \{\varphi \mathbb{U}_i \psi \mid 1 \le i \le n\} \cup \{ \circ (\varphi \mathbb{U}_i \psi) \mid 1 \le i < n\} \cup \operatorname{cl}(\varphi) \cup \operatorname{cl}(\psi).$ Therefore,

$$|\operatorname{cl}(\varphi \mathbb{U}_n \psi)| \le 2n - 1 + |\operatorname{cl}(\varphi)| + |\operatorname{cl}(\psi)|.$$

We consider two different cases:

1. $n = \max\{n, k_{\varphi}, k_{\psi}\}$. Consequently, $2(n - k_{\varphi})|\varphi| \ge 0$ and $2(n - k_{\psi})|\psi| \ge 0$:

$$\begin{aligned} |\mathrm{cl}(\varphi \mathbb{U}_n \psi)| &\leq 2n - 1 + |\mathrm{cl}(\varphi)| + |\mathrm{cl}(\psi)| \\ &\leq 2n - 1 + 2k_{\varphi}|\varphi| + 2k_{\psi}|\psi| \qquad \text{by induction on } \varphi \text{ and } \psi \\ &\leq 2n - 1 + 2k_{\varphi}|\varphi| + 2k_{\psi}|\psi| + 2(n - k_{\varphi})|\varphi| + 2(n - k_{\psi})|\psi| \\ &\leq 2n - 1 + 2k_{\varphi}|\varphi| + 2k_{\psi}|\psi| + 2n|\varphi| - 2k_{\varphi}|\varphi| + 2n|\psi| - 2k_{\psi}|\psi| \\ &\leq 2n - 1 + 2n|\varphi| + 2n|\psi| \\ &\leq 2n - 1 + 2n(|\varphi| + |\psi|) \\ &\leq 2n - 1 + 2n(|\varphi| + |\psi|) \\ &\leq 2n - 1 - 2n + 2n|\varphi \mathbb{U}_n \psi| - 1) \\ &\leq 2n - 1 - 2n + 2n|\varphi \mathbb{U}_n \psi| \\ &\leq 2n|\varphi \mathbb{U}_n \psi|. \end{aligned}$$

2. otherwise, assume that $k_{\psi} = \max\{k_{\varphi}, k_{\psi}\}$. Therefore, $2(k_{\psi} - k_{\varphi})|\varphi| \ge 0$:

$$\begin{aligned} |\mathrm{cl}(\varphi \mathbb{U}_n \psi)| &\leq 2n - 1 + |\mathrm{cl}(\varphi)| + |\mathrm{cl}(\psi)| \\ &\leq 2n - 1 + 2k_{\varphi}|\varphi| + 2k_{\psi}|\psi| \qquad \text{by induction on } \varphi \text{ and } \psi \\ &\leq 2n - 1 + 2k_{\varphi}|\varphi| + 2k_{\psi}|\psi| + 2(k_{\psi} - k_{\varphi})|\varphi| \end{aligned}$$

$$\begin{split} &\leq 2n - 1 + 2k_{\varphi}|\varphi| + 2k_{\psi}|\psi| + 2k_{\psi}|\varphi| - 2k_{\varphi}|\varphi| \\ &\leq 2n - 1 + 2k_{\psi}|\varphi| + 2k_{\psi}|\psi| \\ &\leq 2n - 1 + 2k_{\psi} \left(|\varphi| + |\psi|\right) \\ &\leq 2n - 1 + 2k_{\psi} \left(|\varphi \mathbb{U}_{n}\psi| - 1\right) \\ &\leq 2n - 1 - 2k_{\psi} + 2k_{\psi}|\varphi \mathbb{U}_{n}\psi| \\ &\leq 2k_{\psi}|\varphi \mathbb{U}_{n}\psi|. \qquad \qquad 2n - 1 - 2k_{\psi} < 0 \end{split}$$

• The proof for $\varphi \mathbb{R}_n \psi$, $\varphi \mathbf{S}_n \psi$ and $\varphi \mathbf{T}_n \psi$ follows the same line of reasoning as for $\varphi \mathbb{U}_n \psi$.

QED

Proof of Proposition 14. We proceed by structural induction on μ . We consider only the the metric operators.

• If $\mu = \varphi \mathbb{U}_n \psi$ we proceed by cases. If $\overline{n} = 1$ we conclude that

$$\boldsymbol{m}(k,\mathbf{L}_{\mu}) \stackrel{\eta(\mu)}{=} \boldsymbol{m}(k,\mathbf{L}_{\psi}) \stackrel{IH}{=} \boldsymbol{m}(k,\psi) \stackrel{Prop. 6}{=} \boldsymbol{m}(k,\mu).$$

If $\overline{n} > 1$, let us take $\varphi' = O(\varphi \mathbb{U}_{n-1} \psi)$ to conclude that

$$\boldsymbol{m}(k,\mathbf{L}_{\mu}) \stackrel{\eta(\mu)}{=} \boldsymbol{m}(k,\mathbf{L}_{\psi} \vee (\mathbf{L}_{\varphi} \wedge \mathbf{L}_{\varphi'})) \stackrel{IH}{=} \boldsymbol{m}(k,\psi \vee (\varphi \wedge \varphi')) \stackrel{Prop. 6}{=} \boldsymbol{m}(k,\mu).$$

For the case of ℓ let us take $\varphi' = O(\varphi \mathbb{U}_{\ell} \psi)$ to conclude that

$$\boldsymbol{m}(k,\mathbf{L}_{\mu}) \stackrel{\eta(\mu)}{=} \boldsymbol{m}(k,\mathbf{L}_{\psi} \vee (\mathbf{L}_{\varphi} \wedge \mathbf{L}_{\varphi'})) \stackrel{IH}{=} \boldsymbol{m}(k,\psi \vee (\varphi \wedge \varphi')) \stackrel{Prop. 7}{=} \boldsymbol{m}(k,\mu).$$

• If $\mu = \varphi \mathbb{R}_n \psi$ we proceed by cases. If $\overline{n} = 1$ we use the first formula to conclude that

$$\boldsymbol{m}(k,\mathbf{L}_{\mu}) \stackrel{\eta(\mu)}{=} \boldsymbol{m}(k,\mathbf{L}_{\psi}) \stackrel{IH}{=} \boldsymbol{m}(k,\psi) = \boldsymbol{m}(k,\mu)$$

If $\overline{n} > 1$, let us take $\varphi' = \widehat{O}(\varphi \mathbb{R}_{n-1} \psi)$ to conclude that

$$\boldsymbol{m}(k,\mathbf{L}_{\mu}) \stackrel{\eta(\mu)}{=} \boldsymbol{m}(k,\mathbf{L}_{\psi} \wedge (\mathbf{L}_{\varphi} \vee \mathbf{L}_{\varphi'})) \stackrel{IH}{=} \boldsymbol{m}(k,\psi \wedge (\varphi \vee \varphi')) \stackrel{Prop. 6}{=} \boldsymbol{m}(k,\mu).$$

For the case of ℓ let us take $\varphi' = \widehat{\circ} (\varphi \mathbb{R}_{\ell} \psi)$ to conclude that

$$\boldsymbol{m}(k,\mathbf{L}_{\mu}) \stackrel{\eta(\mu)}{=} \boldsymbol{m}(k,\mathbf{L}_{\psi} \wedge (\mathbf{L}_{\varphi} \vee \mathbf{L}_{\varphi'})) \stackrel{IH}{=} \boldsymbol{m}(k,\psi \wedge (\varphi \vee \varphi')) \stackrel{Prop. 7}{=} \boldsymbol{m}(k,\mu).$$

• If $\mu = \varphi \mathbf{S}_n \psi$ we proceed by cases. If $\overline{n} = 1$ we use the first formula to conclude that

$$\boldsymbol{m}(k,\mathbf{L}_{\mu}) \stackrel{\eta(\mu)}{=} \boldsymbol{m}(k,\mathbf{L}_{\psi}) \stackrel{IH}{=} \boldsymbol{m}(k,\psi) = \boldsymbol{m}(k,\mu)$$

If $\overline{n} > 1$, let us take $\varphi' = \bullet (\varphi \mathbf{S}_{n-1} \psi)$ to conclude that

$$\boldsymbol{m}(k,\mathbf{L}_{\mu}) \stackrel{\eta(\mu)}{=} \boldsymbol{m}(k,\mathbf{L}_{\psi} \vee (\mathbf{L}_{\varphi} \wedge \mathbf{L}_{\varphi'})) \stackrel{IH}{=} \boldsymbol{m}(k,\psi \vee (\varphi \wedge \varphi')) \stackrel{Prop. 6}{=} \boldsymbol{m}(k,\mu).$$

For the case of $\pmb{\ell},$ let us take $\varphi'= \pmb{\bullet}\,(\varphi \pmb{\mathsf{S}}_{\pmb{\ell}}\psi)$ to conclude that

$$\boldsymbol{m}(k,\mathbf{L}_{\mu}) \stackrel{\eta(\mu)}{=} \boldsymbol{m}(k,\mathbf{L}_{\psi} \vee (\mathbf{L}_{\varphi} \wedge \mathbf{L}_{\varphi'})) \stackrel{IH}{=} \boldsymbol{m}(k,\psi \vee (\varphi \wedge \varphi')) \stackrel{Prop. 7}{=} \boldsymbol{m}(k,\mu).$$

• If $\mu = \varphi \mathbf{T}_n \psi$ we proceed by cases. If $\overline{n} = 1$ we use the first formula to conclude that

$$\boldsymbol{m}(k, \mathbf{L}_{\mu}) \stackrel{\eta(\mu)}{=} \boldsymbol{m}(k, \mathbf{L}_{\psi}) \stackrel{IH}{=} \boldsymbol{m}(k, \psi) = \boldsymbol{m}(k, \mu)$$

If $\overline{n} > 1$, let us take $\varphi' = \widehat{\bullet} (\varphi \mathsf{T}_{n-1} \psi)$ to conclude that

$$\boldsymbol{m}(k,\mathbf{L}_{\mu}) \stackrel{\eta(\mu)}{=} \boldsymbol{m}(k,\mathbf{L}_{\psi} \wedge (\mathbf{L}_{\varphi} \vee \mathbf{L}_{\varphi'})) \stackrel{IH}{=} \boldsymbol{m}(k,\psi \wedge (\varphi \vee \varphi')) \stackrel{Prop. 6}{=} \boldsymbol{m}(k,\mu).$$

For the case of ℓ , let us take $\varphi' = \widehat{\bullet} (\varphi \mathsf{T}_{\ell} \psi)$ to conclude that

$$\boldsymbol{m}(k,\mathbf{L}_{\mu}) \stackrel{\eta(\mu)}{=} \boldsymbol{m}(k,\mathbf{L}_{\psi} \wedge (\mathbf{L}_{\varphi} \vee \mathbf{L}_{\varphi'})) \stackrel{IH}{=} \boldsymbol{m}(k,\psi \wedge (\varphi \vee \varphi')) \stackrel{Prop. 7}{=} \boldsymbol{m}(k,\mu).$$

QED

Proof of Theorem 2. Take the MEL_f-trace $\langle \mathbf{H}', \mathbf{T}' \rangle$ whose three valued interpretation m' satisfies:

$$m'(k, \mathbf{L}_{\varphi}) = m(k, \varphi)$$

for any formula φ over \mathcal{A} and for all $i \in [k..\lambda)$. When φ is an atom $a \in \mathcal{A}$ then $\mathbf{m'}(k, a) = \mathbf{m'}(k, \mathbf{L}_a) = \mathbf{m}(k, a)$, which implies that both valuations coincide for atoms, and so, $\langle \mathbf{H'}, \mathbf{T'} \rangle |_{\mathcal{A}} = \langle \mathbf{H}, \mathbf{T} \rangle$. It remains to be shown that $\langle \mathbf{H'}, \mathbf{T'} \rangle \models v(\varphi)$, which is equivalent to

The first satisfaction relation follows directly from the definition of $\langle \mathbf{H}', \mathbf{T}' \rangle$ since $\mathbf{m}'(0, \mathbf{L}_{\varphi}) = 2$ iff $\mathbf{m}(0, \varphi) = 2$ and we had that $\langle \mathbf{H}, \mathbf{T} \rangle$ is a model of φ .

For the second part, we consider the following cases depending on the structure of μ :

- The boolean connectives and temporal formulas $\circ\varphi$, $\bullet\varphi$, $\widehat{\bullet}\varphi$ and $\widehat{\circ}\varphi$ are left to the reader.
- For the formula $\mu = \varphi \mathbb{U}_1 \psi$, note that $\varphi, \psi \in \mathrm{cl}(\mu)$. Let us reason as follows

$$\boldsymbol{m}'(k,\mathbf{L}_{\mu}) = \boldsymbol{m}(k,\mu) = \boldsymbol{m}(k,\varphi \mathbb{U}_{1}\psi) \stackrel{Prop. \ 6}{=} \boldsymbol{m}(k,\psi) = \boldsymbol{m}(k,\mathbf{L}_{\psi}).$$

The cases $\varphi \mathbb{R}_1 \psi$, $\varphi \mathbf{S}_1 \psi$ and $\varphi \mathbf{T}_1 \psi$ are proved in a similar way.

• For the formula $\mu = \varphi \mathbb{U}_n \psi$, with $\overline{n} > 1$, let us take $\varphi' = \circ (\varphi \mathbb{U}_{n-1} \psi)$. Note that, by definition $\varphi, \psi, \varphi' \in cl(\mu)$. Having said that, we reason as follows

$$\boldsymbol{m}'(k, \mathbf{L}_{\mu}) = \boldsymbol{m}(k, \mu) = \boldsymbol{m}(k, \varphi \mathbb{U}_{n} \psi)$$

$$\stackrel{Prop. 6}{=} \boldsymbol{m}(k, \psi \lor (\varphi \land \bigcirc (\varphi \mathbb{U}_{n-1} \psi)))$$

$$= \max\{\boldsymbol{m}(k, \psi), \min(\boldsymbol{m}(k, \varphi), \boldsymbol{m}(k, \bigcirc (\varphi \mathbb{U}_{n-1} \psi)))\}$$

$$= \max\{\boldsymbol{m}(k, \mathbf{L}_{\psi}), \min(\boldsymbol{m}(k, \mathbf{L}_{\varphi}), \boldsymbol{m}(k, \mathbf{L}_{\varphi'}))\}$$

$$= \boldsymbol{m}(k, \mathbf{L}_{\psi} \lor (\mathbf{L}_{\varphi} \land \mathbf{L}_{\varphi'})).$$

• For the formula $\mu = \varphi \mathbb{R}_n \psi$, with $\overline{n} > 1$, let us take $\varphi' = \widehat{\circ} (\varphi \mathbb{R}_{n-1} \psi)$. Note that, by definition $\varphi, \psi, \varphi' \in cl(\mu)$. Having said that, we reason as follows

$$\boldsymbol{m}'(k, \mathbf{L}_{\mu}) = \boldsymbol{m}(k, \mu) = \boldsymbol{m}(k, \varphi \mathbb{R}_{n} \psi)$$

$$\stackrel{Prop. 6}{=} \boldsymbol{m}(k, \psi \land (\varphi \lor \widehat{\circ} (\varphi \mathbb{R}_{n-1} \psi)))$$

$$= \min\{\boldsymbol{m}(k, \psi), \max(\boldsymbol{m}(k, \varphi), \boldsymbol{m}(k, \widehat{\circ} (\varphi \mathbb{R}_{n-1} \psi)))\}$$

$$= \min\{\boldsymbol{m}(k, \mathbf{L}_{\psi}), \max(\boldsymbol{m}(k, \mathbf{L}_{\varphi}), \boldsymbol{m}(k, \mathbf{L}_{\varphi'}))\}$$

$$= \boldsymbol{m}(k, \mathbf{L}_{\psi} \lor (\mathbf{L}_{\varphi} \land \mathbf{L}_{\varphi'})).$$

• For the formula $\mu \varphi \mathbf{S}_n \psi$, with $\overline{n} > 1$, let us take $\varphi' = \mathbf{\bullet} (\varphi \mathbf{S}_{n-1} \psi)$. Note that, by definition $\varphi, \psi, \varphi' \in cl(\mu)$. Having said that, we reason as follows

$$\boldsymbol{m'}(k, \mathbf{L}_{\mu}) = \boldsymbol{m}(k, \mu) = \boldsymbol{m}(k, \varphi \mathbf{S}_{n} \psi)$$

$$\stackrel{Prop. 6}{=} \boldsymbol{m}(k, \psi \lor (\varphi \land \bullet (\varphi \mathbf{S}_{n-1} \psi)))$$

$$= \max\{\boldsymbol{m}(k, \psi), \min(\boldsymbol{m}(k, \varphi), \boldsymbol{m}(k, \bullet (\varphi \mathbf{S}_{n-1} \psi)))\}$$

$$= \max\{\boldsymbol{m}(k, \mathbf{L}_{\psi}), \min(\boldsymbol{m}(k, \mathbf{L}_{\varphi}), \boldsymbol{m}(k, \mathbf{L}_{\varphi'}))\}$$

$$= \boldsymbol{m}(k, \mathbf{L}_{\psi} \lor (\mathbf{L}_{\varphi} \land \mathbf{L}_{\varphi'})).$$

• For the formula $\mu = \varphi \mathbf{T}_n \psi$, with $\overline{n} > 1$, let us take $\varphi' = \widehat{\bullet} (\varphi \mathbf{T}_{n-1} \psi)$. Note that, by definition $\varphi, \psi, \varphi' \in cl(\mu)$. Having said that, we reason as follows

$$\boldsymbol{m'}(k, \mathbf{L}_{\mu}) = \boldsymbol{m}(k, \mu) = \boldsymbol{m}(k, \varphi \mathbf{T}_{n}\psi)$$

$$\stackrel{Prop. 6}{=} \boldsymbol{m}(k, \psi \land (\varphi \lor \widehat{\mathbf{o}} (\varphi \mathbf{T}_{n-1}\psi)))$$

$$= \min\{\boldsymbol{m}(k, \psi), \max(\boldsymbol{m}(k, \varphi), \boldsymbol{m}(k, \widehat{\mathbf{o}} (\varphi \mathbf{T}_{n-1}\psi)))\}$$

$$= \min\{\boldsymbol{m}(k, \mathbf{L}_{\psi}), \max(\boldsymbol{m}(k, \mathbf{L}_{\varphi}), \boldsymbol{m}(k, \mathbf{L}_{\varphi'}))\}$$

$$= \boldsymbol{m}(k, \mathbf{L}_{\psi} \lor (\mathbf{L}_{\varphi} \land \mathbf{L}_{\varphi'})).$$

• For the formula $\mu = \varphi \mathbb{U}_{\ell} \psi$, let us take $\varphi' = 0$ ($\varphi \mathbb{U}_{\ell} \psi$). Note that, by definition $\varphi, \psi, \varphi' \in cl(\mu)$. Having said that, we reason as follows

$$\boldsymbol{m'}(k, \mathbf{L}_{\mu}) = \boldsymbol{m}(k, \mu) = \boldsymbol{m}(k, \varphi \mathbb{U}_{\boldsymbol{\ell}} \psi)$$

$$\stackrel{Prop. 7}{=} \boldsymbol{m}(k, \psi \lor (\varphi \land \circ (\varphi \mathbb{U}_{\boldsymbol{\ell}} \psi)))$$

$$= \max\{\boldsymbol{m}(k, \psi), \min(\boldsymbol{m}(k, \varphi), \boldsymbol{m}(k, \circ (\varphi \mathbb{U}_{\boldsymbol{\ell}} \psi)))\}$$

$$= \max\{\boldsymbol{m}(k, \mathbf{L}_{\psi}), \min(\boldsymbol{m}(k, \mathbf{L}_{\varphi}), \boldsymbol{m}(k, \mathbf{L}_{\varphi'}))\}$$

$$= \boldsymbol{m}(k, \mathbf{L}_{\psi} \lor (\mathbf{L}_{\varphi} \land \mathbf{L}_{\varphi'})).$$

• For the formula $\mu = \varphi \mathbb{R}_{\ell} \psi$, let us take $\varphi' = \widehat{O}(\varphi \mathbb{R}_{\ell} \psi)$. Note that, by definition $\varphi, \psi, \varphi' \in cl(\mu)$. Having said that, we reason as follows

$$\boldsymbol{m'}(k, \mathbf{L}_{\mu}) = \boldsymbol{m}(k, \mu) = \boldsymbol{m}(k, \varphi \mathbb{R}_{\boldsymbol{\ell}} \psi)$$

$$\stackrel{Prop. 7}{=} \boldsymbol{m}(k, \psi \land (\varphi \lor \widehat{\circ} (\varphi \mathbb{R}_{\boldsymbol{\ell}} \psi)))$$

$$= \min\{\boldsymbol{m}(k, \psi), \max(\boldsymbol{m}(k, \varphi), \boldsymbol{m}(k, \widehat{\circ} (\varphi \mathbb{R}_{\boldsymbol{\ell}} \psi)))\}$$

$$= \min\{\boldsymbol{m}(k, \mathbf{L}_{\psi}), \max(\boldsymbol{m}(k, \mathbf{L}_{\varphi}), \boldsymbol{m}(k, \mathbf{L}_{\varphi'}))\}$$

$$= \boldsymbol{m}(k, \mathbf{L}_{\psi} \lor (\mathbf{L}_{\varphi} \land \mathbf{L}_{\varphi'})).$$

• For the formula $\mu = \varphi \mathbf{S}_{\ell} \psi$, let us take $\varphi' = \mathbf{\bullet} (\varphi \mathbf{S}_{\ell} \psi)$. Note that, by definition $\varphi, \psi, \varphi' \in \mathrm{cl}(\mu)$. Having said that, we reason as follows

$$\boldsymbol{m'}(k, \mathbf{L}_{\mu}) = \boldsymbol{m}(k, \mu) = \boldsymbol{m}(k, \varphi \mathbf{S}_{\ell} \psi)$$

$$\stackrel{Prop. 7}{=} \boldsymbol{m}(k, \psi \lor (\varphi \land \bullet (\varphi \mathbf{S}_{\ell} \psi)))$$

$$= \max\{\boldsymbol{m}(k, \psi), \min(\boldsymbol{m}(k, \varphi), \boldsymbol{m}(k, \bullet (\varphi \mathbf{S}_{\ell} \psi)))\}$$

$$= \max\{\boldsymbol{m}(k, \mathbf{L}_{\psi}), \min(\boldsymbol{m}(k, \mathbf{L}_{\varphi}), \boldsymbol{m}(k, \mathbf{L}_{\varphi'}))\}$$

$$= \boldsymbol{m}(k, \mathbf{L}_{\psi} \lor (\mathbf{L}_{\varphi} \land \mathbf{L}_{\varphi'})).$$

• For the formula $\mu = \varphi \mathsf{T}_{\ell} \psi$, let us take $\varphi' = \widehat{\bullet} (\varphi \mathsf{T}_{\ell} \psi)$. Note that, by definition $\varphi, \psi, \varphi' \in \mathrm{cl}(\mu)$. Having said that, we reason as follows

$$\boldsymbol{m'}(k, \mathbf{L}_{\mu}) = \boldsymbol{m}(k, \mu) = \boldsymbol{m}(k, \varphi \mathbf{T}_{\boldsymbol{\ell}} \psi)$$

$$\stackrel{Prop. 7}{=} \boldsymbol{m}(k, \psi \land (\varphi \lor \widehat{\mathbf{0}} (\varphi \mathbf{T}_{\boldsymbol{\ell}} \psi)))$$

$$= \min\{\boldsymbol{m}(k, \psi), \max(\boldsymbol{m}(k, \varphi), \boldsymbol{m}(k, \widehat{\mathbf{0}} (\varphi \mathbf{T}_{\boldsymbol{\ell}} \psi)))\}$$

$$= \min\{\boldsymbol{m}(k, \mathbf{L}_{\psi}), \max(\boldsymbol{m}(k, \mathbf{L}_{\varphi}), \boldsymbol{m}(k, \mathbf{L}_{\varphi'}))\}$$

$$= \boldsymbol{m}(k, \mathbf{L}_{\psi} \lor (\mathbf{L}_{\varphi} \land \mathbf{L}_{\varphi'})).$$

QED